

MedStar Family Choice Prior Authorization and Step Therapy Table

Disclaimer: Medically accepted indications are defined using the following sources: the Food and Drug Administration (FDA), Micromedex, American Hospital Formulary Service (AHFS), United States Pharmacopeia Drug Information for the Healthcare Professional (USP DI), and the Drug Package Insert (PPI).

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
abaloparatide (Tymlos) 3120mcg/1.56ml	<ol style="list-style-type: none"> 1. Prescribed for an approved indication for use: <ul style="list-style-type: none"> • Treatment of postmenopausal women with osteoporosis at high risk for fracture, or patients who have failed or are intolerant to other available osteoporosis therapy. • Treatment to increase bone density in men with osteoporosis at high risk for fracture, or patients who have failed or intolerant to other available osteoporosis therapy. 2. Patient has diagnosis of post-menopausal osteoporosis and is at high risk for bone fracture. 3. Patient is female, age \geq 18 years of age. 4. Patient does not have increased baseline risk for osteosarcoma (e.g., Paget's disease of the bone, bone metastases, or skeletal malignancies). 5. T-score \leq -2.5 based on BMD measurements from the lumbar spine (at least two vertebral bodies), hip (femoral neck, total hip), or radius (one-third radius site) OR <ul style="list-style-type: none"> • History of one of the following resulting from minimal trauma: vertebral compression fracture, fracture of the hip, fracture of the distal radius, fracture of the pelvis, fracture of the proximal humerus. 6. If the criteria in #2 are not met, approval may be granted for patients with both of the following: 	<ol style="list-style-type: none"> 1. Treatment duration has not exceeded a total of 24 months of cumulative use of parathyroid hormone analogs (e.g., Teriparatide, Forteo, Tymlos) during the patient's lifetime. 2. Up to 12 months, not intended to last longer than the final infusion completing 24 months of therapy.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> • BMD T-score between -1 and -2.5 based on BMD measurements from lumbar spine, hip, or radius; AND • ONE of the following FRAX 10-year fracture probabilities: <ul style="list-style-type: none"> ○ Major osteoporotic fracture ≥ 20% ○ Hip fracture ≥ 3% <ol style="list-style-type: none"> 7. Documented trial of teriparatide (Forteo). 8. Documented intolerance, ineffectiveness, contraindication, and/or treatment failure of a minimum trial of 12 weeks of an oral bisphosphonate product. 9. Treatment duration has not exceeded a total of 24 months of cumulative use of parathyroid hormone analogs (e.g., Teriparatide, Forteo, Tymlos) during the patient's lifetime. 10. Approval Duration: up to 12 months 	
adagrasib (Krazati) tablets 200mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • To treat <i>KRAS</i> G12C-mutated locally advanced or metastatic non-small cell lung cancer (NSCLC), as determined by an approved test, in adults who have received at least 1 prior systemic therapy. 2. Test results confirming presence of <i>KRAS</i> G12C mutation in tumor or plasma specimens. 3. Patient has had at least one prior systemic therapy. 4. Medication ordered by an Oncologist. 5. Approval Duration: 12 months. 	<ol style="list-style-type: none"> 1. Confirmation that medication still carries FDA-approval for intended indication. 2. Prescriber has submitted documentation showing periodic monitoring of AST, ALT, alkaline phosphatase, and total bilirubin. 3. No documentation of disease progression or unacceptable toxicity. 4. Approval Duration: 12 months
albuterol inhalers levalbuterol inhalers	<ol style="list-style-type: none"> 1. If patient has exceeded 6 inhalers per 365 days: Note: this applies to any combination of albuterol MDIs and levalbuterol MDIs. <ul style="list-style-type: none"> • Provider must show that patient has been prescribed appropriate controller therapy for 	

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>indication (asthma, COPD).</p> <ul style="list-style-type: none"> • Provider must provide documentation of treatment plan and patient follow-up that will occur. • Patient must be referred for follow up with MFC Case Management. • Approval for asthma indication is for on <p>2. Approval for asthma indication is for one fill, one month only.</p> <p>3. Approval for COPD may be longer depending upon documented COPD severity, concurrent therapy, compliance with COPD maintenance medication regimen, and oversight by a pulmonologist.</p>	
alectinib (Alecensa) capsule 150mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Treatment of patients with anaplastic lymphoma kinase (ALK)- positive metastatic non-small cell lung cancer (NSCLC) as detected by an FDA-approved test. 2. Patient ≥ 18 years of age. 3. Patient has advanced or metastatic disease. 4. Patient has anaplastic lymphoma kinase (ALK)- positive disease as detected by an approved test. 5. Medication ordered by an Oncologist. 6. Maximum Approval Duration: 12 months. 	<ol style="list-style-type: none"> 1. No documentation of disease progression or unacceptable toxicity. 2. Authorization Duration: 12 months.
alosetron (Lotronex) 0.5 mg, 1 mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • For females with severe diarrhea-predominant irritable bowel syndrome (IBS), including one or more of the following criteria: <ul style="list-style-type: none"> • Frequent and severe abdominal pain/discomfort • Frequent bowel urgency or fecal incontinence • Disability or restriction of daily activities due to 	<ol style="list-style-type: none"> 1. Documentation of positive clinical response to therapy. 2. Authorization Duration: 12 months.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p style="text-align: center;">IBS</p> <ol style="list-style-type: none"> 2. Chronic IBS symptoms lasting at least 6 months. 3. Gastrointestinal tract abnormalities have been ruled out 4. There has been an inadequate response to conventional therapy (e.g. loperamide, antispasmodics). 5. The patient does not have a history of any of the following conditions: <ul style="list-style-type: none"> • Chronic or severe constipation or sequelae from constipation • Intestinal obstruction, stricture, toxic megacolon, gastrointestinal perforation, and/or adhesions • Ischemic colitis • Impaired intestinal circulation, thrombophlebitis, or hypercoagulable state • Crohn’s disease or ulcerative colitis • Diverticulitis • Severe hepatic impairment 6. Dose is limited to 2 tablets per day. 7. Initial authorization is for 6 months. 	
armodafinil (Nuvigil) tablets 50mg, 150mg 200mg, 250mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • To improve wakefulness in adult patients with excessive sleepiness associated with obstructive sleep apnea, narcolepsy, or shift work disorder. 2. Medication ordered by a Neurologist or certified sleep specialist. 	
avatrombopag (Doptelet) tablets 20mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • thrombocytopenia in adult patients with chronic liver disease who are scheduled to undergo a procedure. 	When prescribed for thrombocytopenia in chronic liver disease with procedure scheduled:

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> • thrombocytopenia in adult patients with chronic immune thrombocytopenia who have had an insufficient response to a previous treatment. <ol style="list-style-type: none"> 2. Patient age \geq 18 years. 3. A recent (less than 1 month old) platelet count must be supplied with documentation submitted. 4. Medication ordered by a Hematologist. <p>When prescribed for thrombocytopenia in patients with chronic liver disease-associated thrombocytopenia scheduled to undergo a procedure:</p> <ul style="list-style-type: none"> • Approval limited to 15 tablets per treatment course. • Approval Duration: one month. <p>When prescribed to patients with chronic immune thrombocytopenia with insufficient response to previous treatment:</p> <ul style="list-style-type: none"> • Diagnosis of chronic immune thrombocytopenia (ITP). • Patient experienced insufficient response to a previous treatment (e.g., corticosteroids, immunoglobulins, thrombopoietin receptor agonists, splenectomy). • Approval duration: 12 months. 	<ul style="list-style-type: none"> • must meet initial use criteria for each request. • Maximum approval duration: 1 month • Maximum of 15 tablets per treatment. <p>When prescribed to patients with chronic immune thrombocytopenia with insufficient response to previous treatment:</p> <ul style="list-style-type: none"> • Documented positive response to treatment. • Approval Duration: 12 months.
axicabtagene ciloleucel (Yescarta) Injection	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • treatment of adult patients with large B- cell lymphoma that is refractory to first-line chemotherapy. 2. The treatment facility that dispenses and administers Yescarta is enrolled and complies with the Risk Evaluation and Mitigation Strategy; AND 3. Patient age \geq 18 years. 4. Patient has received prior treatment with first-line 	<p>Not applicable. Maximum approval, one treatment course per lifetime.</p>

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	chemo-immunotherapy and has any of the following B-cell lymphoma sub-types: <ul style="list-style-type: none"> • Diffuse large B-cell lymphoma (DLBCL) • Primary mediastinal large B-cell lymphoma • High-grade B-cell lymphomas • HIV-related B-cell lymphomas • Monomorphic post-transplant lymphoproliferative B-cell type disorder 5. Patient does not have ANY of the following: <ul style="list-style-type: none"> • Primary CNS lymphoma • Previous treatment with Yescarta or other CD 19-directed chimeric antigen receptor (CAR) T-cell therapy. • ECOG performance status ≥ 3 (patient is not ambulatory, capable of self-care, or confined to bed or chair more than 50% of waking hours). • Inadequate or unstable kidney, liver, pulmonary, or cardiac function. • Active hepatitis B, active hepatitis C, or clinically active systemic infection. 6. Medication ordered by an Oncologist. 7. Approval duration: 3 months.	
azacitadine (Onureg) tablets 200mg, 300mg	1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Continued treatment of adult patients with acute myeloid leukemia who achieved first complete remission or complete remission with incomplete blood count recovery following intensive induction chemotherapy and are not able to complete intensive curative therapy. 2. Patient is not able to complete intensive curative	1. Patient does not show evidence of progressive disease while on Onureg therapy. 2. Approval duration: 12 months.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	therapy (i.e. transplant-ineligible). 3. Medication ordered by an Oncologist.	
azelaic acid (Finacea) gel 15% STEP THERAPY	1. Ordered for Acne Vulgaris in adults. 2. Patient has had an adequate trial (30 days) of at least two types of formulary, topical acne products. Two types meaning, two different active ingredients. <ul style="list-style-type: none"> Acceptable formulary precursor ingredients include: adapalene, benzoyl peroxide, benzoyl peroxide-erythromycin combination products, clindamycin, clindamycin-benzoyl peroxide combination products, erythromycin, tretinoin. 3. If patient's claims data supports the completion of the step-therapy, the claim will adjudicate without manual review. 4. Approval Duration: 12 months.	
baricitinib (Olumiant) tablets 2 mg, 4 mg (1 mg non-formulary)	1. Ordered for an approved indication for use: <ul style="list-style-type: none"> Treatment of patients with severe alopecia areata Treatment of patients with moderately to severely active rheumatoid arthritis who have had an inadequate response to one or more TNF blockers. 2. Patient age ≥ 18 years. Alopecia Areata: <ul style="list-style-type: none"> Diagnosis of moderate to severe alopecia areata, with a current episode of alopecia areata with at least 50% scalp hair loss, AND Other causes of hair loss have been ruled out (e.g., androgenetic alopecia, cicatricial alopecias, secondary syphilis, tinea capitis, triangular alopecia, and trichotillomania); AND Patient is not receiving Olumiant concurrent with ANY of the following: 	Renewal Criteria applies to both Alopecia Areata and Rheumatoid Arthritis indications: 1. Documentation of positive clinical response to Olumiant therapy; AND 2. Patient is not receiving Olumiant concurrent with ANY of the following: <ul style="list-style-type: none"> A targeted immunomodulator e.g. Adalimumab, Cimzia, Enbrel, Simponi, Orencia, Xeljanz, Rinvoq or Litfulo OR A potent immunosuppressant (e.g. azathioprine or cyclosporine). 3. Approval Duration: 12 months.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> ○ A targeted immunomodulator e.g., adalimumab, Cimzia, Enbrel, Simponi, Orencia, Xeljanz, Rinvoq or Litfulo OR ○ A potent immunosuppressant (e.g., azathioprine or cyclosporine). ● Prescribed by, or in consultation with a Dermatologist. ● Limitations: 1 tablet per day (i.e. if 4 mg dose needed, use 4 mg tablets rather than 2 x 2 mg tablets). ● Approval Duration: 12 months <p><u>Rheumatoid Arthritis (RA):</u></p> <ul style="list-style-type: none"> ● Diagnosis of moderately to severely active RA; AND ● <u>One</u> of the following: <ul style="list-style-type: none"> ○ History of failure to a 3-month trial of one non-biologic disease modifying anti-rheumatic drug (DMARD) (e.g. methotrexate, leflunomide, sulfasalazine, hydroxychloroquine) at maximally indicated doses, unless contraindicated or clinically significant adverse effects are experienced (document drug, date, and duration of trial), OR ○ Patient has previously been treated with a targeted immunomodulator FDA-approved for the treatment of RA as documented by claims history or medical records that include drug, date and duration of therapy. (e.g., adalimumab, Enbrel, Cimzia, Simponi, Orencia, Xeljanz, Rinvoq); AND ● <u>One</u> of the following: 	

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> ○ History of failure, contraindication or intolerance to at least two preferred products with documentation of drug, date, and duration of trial: adalimumab, Enbrel, Cimzia, Simponi, Rinvoq, Xeljanz; OR ○ Patient is currently on Olumiant therapy as documented by claims history or submission of medical records AND the patient has not received a manufacturer supplied sample at no cost as a means to establish themselves as a current user of Olumiant; AND ● Patient is not receiving Olumiant concurrent with ANY of the following: <ul style="list-style-type: none"> ○ A targeted immunomodulator e.g., adalimumab, Cimzia, Enbrel, Simponi, Orencia, Xeljanz, Rinvoq or Litfulo OR ○ A potent immunosuppressant (e.g., azathioprine or cyclosporine). ● Prescribed by or in consultation with a Rheumatologist. ● Dose is limited to one, 2 mg per day. ● Approval Duration: 12 months. 	
bedaquiline (Sirturo) tablets 20mg, 100mg	1. Ordered for an approved indication for use: <ul style="list-style-type: none"> ● as part of combination therapy in adult and pediatric patients ≥ 5 years of age and weighing at least 15 kg with pulmonary multi-drug resistant tuberculosis (MDR-TB). [Reserved for use when an effective treatment regimen cannot otherwise be provided. 	

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	2. Medication ordered by an infectious disease specialist 3. Approval duration: 24 weeks	
belimumab (Benlysta) Inj 200mg/ml	1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • patients ≥ 5 years of age with active systemic lupus erythematosus (SLE) who are receiving standard therapy. • Patients ≥ 18 years of age with active lupus nephritis who are receiving standard therapy. 2. Not prescribed for patients with severe active central nervous system lupus as use of Benlysta is not recommended for those patients. 3. Must be currently taking OR has tried and failed or had intolerance/contraindication to at least one standard therapy for SLE (e.g., corticosteroids, antimalarials, NSAIDS or immunosuppressives) or lupus nephritis (e.g., corticosteroids, mycophenolate, cyclophosphamide, azathioprine) 4. Prescriber attestation that all baseline evaluations have been done and no contraindications to use are present including counseling/assessment of recent live vaccine use and depression/suicide risk. 5. Prescriber attests that subsequent appropriate evaluation and monitoring will be done based on the package insert. 6. Patient is not receiving Benlysta in combination with ANY of the following: <ul style="list-style-type: none"> • Targeted Immunomodulator (e.g., Adalimumab, etanercept, certolizumab, anakinra) 	1. All initial criteria continue to be met. 2. Documentation demonstrating clinical benefit and tolerance. 3. Patient is not receiving Benlysta in combination with ANY of the following: <ul style="list-style-type: none"> • Targeted Immunomodulator (e.g., Adalimumab, etanercept, certolizumab, anakinra) • Lupkynis (voclosporin) • Saphnelo (anifrolumab) 4. Approval Duration: 12 months.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> • Lupkynis (voclosporin) • Saphnelo (anifrolumab) <p>7. Prescribed by an immunologist, nephrologist, rheumatologist, or provider experienced in the treatment of SLE or lupus nephritis.</p> <p>8. Approval Duration: 12 months</p>	
belumosudil (Rezurock) tablets 200mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • treatment of adult and pediatric patients 12 years and older with chronic graft-versus-host disease (chronic GVHD) after failure of at least two prior lines of systemic therapy. 2. Member must have tried and failed, have intolerance or medical contraindication to at least three of these medications: cyclosporine, methotrexate, mycophenolate, sirolimus, and glucocorticoids. 3. Patient age ≥ 12 years. 4. Provider attestation: Drug specific baseline evaluation and monitoring completed (CBC/CMP including total bilirubin, AST, ALT). Patient is not pregnant and is using effective contraception, concurrent use of CYP3A inducers and proton pump inhibitors is contraindicated. 5. Life expectancy is > 6 months. 6. Quantity limited to 30 tablets per 30 days. 7. Approval duration: 6 months 	<ol style="list-style-type: none"> 1. Prescriber attestation of continued clinical benefit. 2. Approval Duration: 6 months.
benralizumab (Fasenra) Pen 10mg/0.5ml, 30mg/ml	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • add-on maintenance treatment of patients ≥ 12 years of age with severe asthma and with an eosinophilic phenotype. 2. Diagnosis of severe, uncontrolled asthma as defined by at least ONE of the following: <ul style="list-style-type: none"> • Poor symptom control (e.g., Asthma Control 	<ol style="list-style-type: none"> 1. Documentation of positive clinical response to Fasenra therapy as demonstrated by at least one of the following: <ul style="list-style-type: none"> • Reduction in frequency of exacerbations • Decreased utilization of rescue

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>Questionnaire (ACQ) score consistently greater than 1.5 or Asthma Control Test (ACT) score consistently less than 20).</p> <ul style="list-style-type: none"> • Two or more bursts of systemic corticosteroids for at least 3 days each in previous 12 months. • Asthma-related emergency treatment (ER visit, hospital admission, or unscheduled OV for nebulizer or emergency treatment). • Airflow limitation (e.g., after appropriate bronchodilator withhold forced expiratory volume in 1 second (FEV1) less than 80% predicted. • Patient is currently dependent on oral corticosteroids for the treatment of asthma. <p>3. Submission of medical records documenting one of the following:</p> <ul style="list-style-type: none"> • Asthma is eosinophilic phenotype as defined by baseline (pre-benralizumab treatment) peripheral blood eosinophil level \geq 150 cells/uL within the past 6 weeks; OR • Patient is currently dependent on maintenance therapy with oral corticosteroids for the treatment of asthma. <p>4. Fasenra will be used in combination with ONE of the following:</p> <ul style="list-style-type: none"> • One high-dose combination inhaled corticosteroid (ICS/LABA); OR • Combination therapy with BOTH one high dose inhaled corticosteroid and one additional asthma controller medication. <p>5. Patient is not receiving treatment in combination with ANY of the following:</p> <ul style="list-style-type: none"> • Anti-interleukin-5 therapy (e.g., Cinqair 	<p>medications</p> <ul style="list-style-type: none"> • Increase in percent predicted FEV1 from pretreatment baseline. • Reduction in severity or frequency of asthma-related symptoms • Reduction in oral corticosteroid requirements. <p>2. Used in combination with inhaled corticosteroid (ICS)-containing controller medication.</p> <p>3. Patient is not receiving treatment in combination with ANY of the following:</p> <ul style="list-style-type: none"> • Anti-interleukin-5 therapy (e.g., Cinqair (reslizumab), Nucala (mepolizumab). • Anti-IgE therapy (e.g., Xolair (omalizumab). • Anti-interleukin-4 therapy (e.g., Dupixent (dupilumab). • Thymic stromal lymphopoietin (TSLP) inhibitor (e.g., Tezpire (Tezepelumab)). <p>4. Approval Duration: 12 months.</p>

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>(reslizumab), Nucala (mepolizumab)).</p> <ul style="list-style-type: none"> • Anti-IgE therapy (e.g., Xolair (omalizumab). • Anti-interleukin-4 therapy (e.g., Dupixent (dupilumab). • Thymic stromal lymphopoietin (TSLP) inhibitor (e.g., Tezspire (Tezepelumab)). <p>6. Medication ordered by a Pulmonologist, Immunologist, or Allergist.</p> <p>7. Approval Duration: 12 months.</p>	
beremagene geperpavec-svdt (Vyjuvek) topical gel 5×10 ⁹ PFU/mL	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Treatment of wounds in patients with diagnosis of dystrophic epidermolysis bullosa (DEB). 2. Patient age ≥ 6 months. 3. Submission of medical records (e.g., chart notes, laboratory values) confirming a mutation in the collagen type VII alpha 1 chain (COL7A1) gene. 4. Patient has at least one recurrent or chronic open wound that meets all of the following: adequate granulation tissue, excellent vascularization AND no evidence of active wound infection. 5. No evidence or history of squamous cell carcinoma. 6. Prescribed by, or in consultation with, a dermatologist with expertise in the treatment of DEB. 7. Dosing is in accordance with FDA approved labeling. Initial authorization limited to no more than 6 months and no more than 26 doses. 	<ol style="list-style-type: none"> 1. Patient has previously been treated with Vyjuvek therapy. 2. Patient had a positive clinical response to Vyjuvek therapy (e.g., decrease in wound size, increase in granulation tissue, complete wound closure). <ol style="list-style-type: none"> 1. Wound(s) being treated to meet all the following criteria: <ul style="list-style-type: none"> • Adequate granulation tissue • Excellent vascularization • No evidence of active wound infection • No evidence or history of squamous cell carcinoma • Dosing is in accordance with FDA approved labeling. <p>Reauthorization limited to no more than 6 months and no more than 26 doses.</p>
berotralstat (Orladeyo) capsules 110mg, 150mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • prophylaxis to prevent attacks of hereditary angioedema (HAE) in adults and pediatric patients 12 years and older. 	<ol style="list-style-type: none"> 1. Member meets the criteria for initial approval.

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ol style="list-style-type: none"> 2. Patient age \geq 12 years. 3. C1 inhibitor (C1-INH) antigenic level below the lower limit of normal as defined by the laboratory performing the test OR Normal X1-inh antigenic level and a low C1-INH functional level (functional C1-INH less than 50% or C1-INH functional level below the lower limit of normal as defined by the laboratory performing the test. 4. Prescriber attestation that all baseline evaluations have been done, prophylactic therapy is medically necessary, and no contraindications to use. 5. Not used in combination with other approved products indicated for prophylaxis against HAE attacks (i.e., Cinryze, Haegarda, Takhzyro). 6. History of failure to BOTH of the following (document date of trial and list reason for therapeutic failure) Haegarda AND Takhzyro. Alternatively- could be failure of one of the following: Haegarda, Cinryze, Takhzyro. 7. Quantity limit of 1 capsule per day 8. Prescriber is a hematologist, immunologist, or allergist. <p>Initial approval duration: 3 months</p>	<ol style="list-style-type: none"> 2. Member has experienced a significant reduction in frequency of attacks (\geq 50%) since starting treatment. 3. Member has reduced the use of medications to treat acute attacks since starting treatment. 4. Prescriber attests that patient has had an annual evaluation for the continued need for long-term prophylaxis therapy AND 5. Prescriber attests a recent review of patient's current medication has been completed and there is no concomitant use of P-gp inducers (e.g. rifampin, St John's wort), and dose adjustment has been made based on labeled recommendations for drug interactions if applicable. 6. Approval Duration: 3 months.
bosutinib (Bosulif) tablets 100mg, 500mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Newly diagnosed chronic phase Ph+ chronic myelogenous leukemia (CML). • Chronic, accelerated, or blast phase Ph+ CML with resistance or intolerance to prior therapy. 2. Medication ordered by an Oncologist. 3. Authorization Duration: 12 months. 	<ol style="list-style-type: none"> 1. Patient does not show evidence of disease progression while on Bosulif therapy. 2. Approval Duration: 12 months.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
brigatinib (Alunbrig) tablets 30mg, 90mg, 180mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • The treatment of adult patients with anaplastic lymphoma kinase (ALK)-positive metastatic non-small cell lung cancer (NSCLC) as detected by an FDA-approved test. • Ordered for treatment when the indication has been recognized by the National Comprehensive Cancer Network (NCCN) Drugs and Biologics Compendium with a category of Evidence and Consensus of 1, 2A, or 2B. <p><u>Non-Small Cell Lung Cancer (NSCLC):</u></p> <ul style="list-style-type: none"> • The tumor is anaplastic lymphoma kinase (ALK)-positive. • The cancer is either: metastatic, recurrent, or advanced. <p><u>Soft Tissue Sarcoma/Uterine Neoplasms:</u></p> <ul style="list-style-type: none"> • Diagnosis of inflammatory myofibroblastic tumor (IMT); and • Presence of ALK translocation. <p><u>Histiocytic Neoplasms:</u></p> <ul style="list-style-type: none"> • Diagnosis of symptomatic Erdheim-Chester Disease; and • Used as targeted therapy ALK-fusion; and • Disease is either relapsed or refractory. <p><u>Central Nervous System (CNS) Cancers:</u></p> <ul style="list-style-type: none"> • Diagnosis of metastatic brain cancer from NSCC; and • Tumor is ALK-positive. 2. Medication ordered by or in consultation with an Oncologist. 3. Approval Duration: 12 months. 	<ol style="list-style-type: none"> 1. Patient does not show evidence of progressive disease while on Alunbrig therapy. 2. Approval Duration: 12 months.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
<p>buprenorphine products for chronic pain (Belbuca, Butrans)</p> <p>buccal film (Belbuca) 75mcg, 150mcg, 300mcg, 450mcg, 600mcg, 750mcg, 900mcg</p> <p>topical patches (Butrans generic) 5mcg/hr, 7.5mcg/hr, 10mcg/hr, 15mcg/hr, 20mcg/hr</p>	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • The management of pain severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate. 2. Requested dose is appropriate based on patient's opioid status: <ul style="list-style-type: none"> • Opioid-naïve or non-opioid tolerant patients <ul style="list-style-type: none"> ○ Buccal film – 75mcg once daily or Q12 for at least 4 days before titration ○ Topical patch – 5mcg/hr once every 7days, wait at least 72 hours before titration • Opioid-tolerant patients – convert based on patient's current opioid regimen <ul style="list-style-type: none"> ○ Buccal film – patients currently receiving >160 MME/day of other opioid may not receive adequate analgesic effect from buccal film at max doses; consider alternative opioid ○ Topical patch – patients currently receiving >80 MME/day of other opioid may not receive adequate analgesic effect from topical patch at max doses; consider alternative opioid 	<ul style="list-style-type: none"> • All long-acting opioids require Prior Authorization (PA). The PA form can be accessed using the following link: OPIOID PRIOR AUTH FORM-DC <p>Limitations of Use: Because of the risks of addiction, abuse, and misuse with opioids, even at recommended doses, and greater risk of overdose and death with extended-release opioid formulations, reserve for use in patients for whom alternative treatment options (e.g., non-opioid analgesics or immediate-release opioids) are ineffective, not tolerated, or would be otherwise inadequate to provide sufficient management of pain.</p>
<p>buprenorphine/naloxone (Suboxone) sublingual film</p> <p>2-0.5, 4-1, 8-2, 12-3 mg</p> <p>*Note: generic buprenorphine/naloxone is</p>	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: 2. Previous trial of generic buprenorphine/naloxone sublingual film medication. 3. Prescriber attestation that the patient is unable to use the generic product and has a medical need to use BRAND Suboxone. 4. Approval Duration: 12 months 	<ol style="list-style-type: none"> 1. Patient continues to meet initial approval criteria.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
formulary-preferred and available without prior authorization.		
c1 esterase Inhibitor [Human] Cinryze solution 500 unit Haegarda injection solution 2000unit, 3000unit	<ol style="list-style-type: none"> 5. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Routine prophylaxis to prevent Hereditary Angioedema attacks in patients \geq 6 years of age. 6. Cinryze will be considered for coverage when ALL of the criteria below are met and confirmed with medical documentation. <ul style="list-style-type: none"> • Diagnosis of hereditary angioedema (HAE) confirmed by one of the following: <ul style="list-style-type: none"> ○ Confirmed monoallelic mutation known to cause HAE in either the SERPING1 or F12 gene: OR ○ A C4 level below the lower limit of normal and either C1 inhibitor (C1-INH) antigenic level below the lower limit of normal or C1-INH functional level below the lower limit of normal; AND • Used for prophylaxis of acute HAE attacks: AND • Patient is at least 6 years of age; AND • Patient has experienced the following: <ul style="list-style-type: none"> ○ History of recurrent laryngeal attacks; OR ○ \geq 2 severe episodes/month (ex. debilitating GI or cutaneous effects); OR ○ \geq 5 days/month of debilitating symptoms; AND • Prescribed by an allergist, immunologist, hematologist, or other appropriate specialist; AND • Medications known to cause angioedema (ex. ACE-Inhibitors, estrogens, angiotensin II receptor blockers) have been evaluated and discontinued 	<ol style="list-style-type: none"> 2. All of the criteria for initial therapy must be met; AND 3. Provider attests to a positive clinical response. 4. Continuing therapy will be authorized for 3 months.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>when appropriate.</p> <p>7. Length of Authorization: 3 months when criteria are met.</p>	
<p>cabotegravir (Apretude) extended-release intramuscular injection 600mg/3mL</p>	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • At-risk adults and adolescents weighing at least 35 kg for PrEP to reduce the risk of sexually acquired HIV-1 infection. 2. Individuals must have a negative HIV-1 test prior to initiating APRETUDE and prior to each injection. 3. Provider confirms that the patient will be tested for HIV-1 infection with each subsequent injection; AND 4. Patient is not an appropriate candidate for oral PrEP (e.g., difficulty with adherence to prior oral PrEP, significant renal disease); AND 5. Provider attests that patient demonstrates treatment readiness by BOTH of the following: 6. Patient understands the risks of missed doses. 7. Patient has ability to adhere to the required every 2 months injection and testing appointments. 8. Dosing is in accordance with FDA-approved labeling. 9. Approval Duration: 2 months 	<ol style="list-style-type: none"> 1. Patient has previously received treatment with Apretude 2. Patient has a negative HIV-1 test 3. Provider confirms that the patient will be tested for HIV-1 with each subsequent injection; and 4. Dosing is in accordance with FDA-approved labeling. 5. Approval Duration: 2 months
<p>cabozantinib (Cabometyx) tablets 20mg, 40mg, 60mg</p>	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Patients with advanced renal cell carcinoma (RCC) • Patients with advanced renal cell carcinoma, as a first-line treatment in combination with nivolumab • Patients with hepatocellular carcinoma (HCC) who have been previously treated with sorafenib • Adult and pediatric patients ≥ 12 years of age with locally advanced or metastatic differentiated thyroid cancer (DTC) that has progressed following prior VEGFR-targeted therapy and who 	

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
caplacizumab-yhdp (Cablivi) kit 11mg	<p style="text-align: center;">are radioactive iodine-refractory or ineligible</p> <ol style="list-style-type: none"> 2. Medication ordered by an Oncologist. <ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Treatment of adult patients with acquired thrombotic thrombocytopenic purpura (aTTP), in combination with plasma exchange and immunosuppressive therapy. 2. Medication ordered by hematologist. 3. The patient received the requested medication with plasma exchange. 4. Cablivi will be given in combination with immunosuppressive therapy. 5. The patient will not receive Cablivi beyond 30 days from the cessation of plasma exchange unless the patient has documented, persistent aTTP. 6. The patient has not experienced more than 2 recurrences of aTTP while on the requested medication. (A recurrence is when the patient needs to reinitiate plasma exchange, a 28-day extension of therapy is not considered a recurrence.) 7. Approval is for 30 days only. 	<ol style="list-style-type: none"> 1. A request for continuation of therapy is for extension of therapy after the initial course of Cablivi. The initial course is treatment with Cablivi during and 30 days after plasma exchange. 2. The patient has either of the following documented signs of persistent, underlying aTTP: <ul style="list-style-type: none"> • ADAMTS13 activity level of < 10%, OR • All of the following: <ul style="list-style-type: none"> ○ Microangiopathic hemolytic anemia (MAHA) documented by the presence of schistocytes on peripheral smear ○ Thrombocytopenia and ○ Elevated lactate dehydrogenase (LDH) level 3. Cablivi will be given in combination with immunosuppressive therapy. 4. The patient has not received a prior 28-day extension of therapy after the initial course of Cablivi. 5. The patient has not experienced more than 2 recurrences of aTTP while on Cablivi. 6. Approval duration: 30 days.
capmatinib (Tabrecta) tablets 150mg, 200mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • treatment of adults with metastatic NSCLC whose 	

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>tumors have a mutation that leads to mesenchymal-epithelial transition (MET) exon 14 skipping as detected by an approved test.</p> <p>2. Medication ordered by an oncologist.</p>	
<p>cariprazine (Vraylar) capsules</p> <p>1.5mg, 3mg, 4.5mg, 6mg Therapy pack – 1.5mg & 3mg</p>	<p>1. Ordered for an approved indication for use:</p> <ul style="list-style-type: none"> • Bipolar disorder • Major depressive disorder, as adjunctive therapy with an antidepressant medication • Schizophrenia <p>2. Patient is ≥18 years</p> <p>3. Patient has completed adequate 12-week trials of at least 3 other antipsychotic medications, or has a contraindication to using other options.</p> <p>4. Medication ordered by a psychiatrist or behavioral health specialist.</p> <p>5. Approval duration: 12 months</p>	
<p>casimersen (Amondys 45) injection 100mg/2ml</p>	<p>1. Ordered for an approved indication for use:</p> <ul style="list-style-type: none"> • Treatment of Duchenne muscular dystrophy (DMD) in patients who have a confirmed mutation of the DMD gene that is amenable to exon 45 skipping. <p>2. Confirmed diagnosis of DMD with genetic confirmation of the DMD gene that is amenable to exon 45 skipping.</p> <p>3. Provider attestation of baseline and subsequent evaluation and monitoring as appropriate such as hypersensitivity reactions and renal function.</p> <p>4. Be on a stable dose of corticosteroid for ≥ 24 weeks.</p> <p>5. Not ventilator dependent</p> <p>6. Not receiving other RNA antisense therapy or gene therapy for DMD.</p>	<ul style="list-style-type: none"> • This indication is approved under accelerated approval based on an increase in dystrophin production in skeletal muscle observed in patients treated with AMONDYS 45. Continued approval for this indication may be contingent upon verification of a clinical benefit in confirmatory trials. • Duration of approval is limited to 6 months. <p>Renewal Criteria:</p> <ul style="list-style-type: none"> • Not receiving other antisense therapy or gene therapy. • Not ventilator dependent.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	7. Maximum dose 30 mg/kg/dose once weekly 8. Prescribed by or in consultation with a pediatric neurologist with expertise in DMD.	<ul style="list-style-type: none"> • Provider attestation of continued benefit without ADE • Max dose 30 mg/kg/dose/week • Duration: 6 months
chlordiazepoxide (Librium) capsules 5mg, 10mg, 25mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ol style="list-style-type: none"> a) alcohol withdrawal syndrome treatment in adults b) management of anxiety disorders 2. No concurrent use of other benzodiazepines 3. If ordered for alcohol withdrawal syndrome: <ol style="list-style-type: none"> a) Current CIWA-AR score 10-15, indicating mild withdrawal symptoms. <ol style="list-style-type: none"> i) Scores >15 – contraindication to ambulatory management ii) Scores <10 – gabapentin is preferred as first-line therapy over benzodiazepines to manage very mild withdrawal symptoms b) No prior history of withdrawal delirium (delirium tremens) or withdrawal seizures c) Confirmation of negative pregnancy status for female patients of reproductive age d) No presence of complex comorbidities or psychiatric comorbidities that can increase the risk of developing severe alcohol withdrawal syndrome: <ol style="list-style-type: none"> i) Heart failure class II or higher ii) Decompensated cirrhosis iii) COPD with oxygen dependence iv) CKD stage IV or higher v) Epilepsy or seizure history vi) Recent head injury with loss of consciousness or intracranial hemorrhage vii) Unstable or active psychiatric illness causing 	Renewal criteria: <ul style="list-style-type: none"> • Alcohol withdrawal syndrome requires a new authorization for each requested treatment course. <ul style="list-style-type: none"> ○ Consider referring to case management if a patient is requested for repeated treatments • Anxiety disorder <ul style="list-style-type: none"> ○ Confirmation of clinically significant improvement in symptoms ○ Duration: 12 months once maintenance dose is established

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>active psychosis, mania, depression , or suicidal ideation</p> <p>viii) Febrile illness</p> <p>ix) Benzodiazepine use disorder</p> <p>e) Evaluation confirming patient is an appropriate candidate for ambulatory management – able to self-monitor symptoms, take medications as directed, attend follow-up visits as instructed, etc.</p> <p>4. If ordered for anxiety disorder management:</p> <p>a) No history of substance use disorder, misuse of medications, or depression</p> <p>b) Established contraindication/failure to using diazepam or clonazepam for this indication</p> <p>c) Adequate 8-12 week trial at a therapeutic dose of a serotonin reuptake inhibitor (SSRIs, SNRIs), or contraindication to use</p> <p>5. Initial approval Duration:</p> <p>a) Alcohol withdrawal syndrome treatment – only approve for the requested treatment regimen; max of #30 for a 4-day supply (25 mg capsules)</p> <p>b) Anxiety disorder management – 8 weeks</p>	
<p>crisaborole (Eucrisa) ointment 2%</p> <p>STEP THERAPY</p>	<p>1. Ordered for an approved indication for use:</p> <ul style="list-style-type: none"> Topical treatment of mild-to-moderate atopic dermatitis in adult and pediatric patients ≥ 3 months of age. <p>2. Step Therapy: Unless patient age <2 years of age. First must have completed a trial of, or have intolerance or contraindication to use BOTH:</p> <ul style="list-style-type: none"> At least one medium- or high-potency topical steroid AND A six-week trial of topical tacrolimus OR 	<p>1. Patient has achieved or maintained a positive clinical response as evidenced by improvement or resolution of any of the following:</p> <ul style="list-style-type: none"> Erythema, edema, xerosis, erosions, excoriations, oozing and crusting, lichenification or pruritus. <p>2. Approval Duration: 12 months.</p>

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>pimecrolimus, OR</p> <ul style="list-style-type: none"> • A four-week trial of Zoryve. <p>3. Quantity limited to 60 grams for a 30 day-supply or 180 grams for a 90 day supply. Additional quantity requests may be granted if the affected area is greater than 5% of body surface area (BSA).</p> <p>4. Approval duration: 12 months</p>	
<p>crizotinib (Xalkori) capsule 200mg, 250mg</p>	<p>1. Ordered for an approved indication for use:</p> <ul style="list-style-type: none"> • the treatment of patients with metastatic non-small cell lung cancer (NSCLC) whose tumors are anaplastic lymphoma kinase (ALK) or ROS1-positive as detected by an FDA-approved test. • pediatric patients 1 year of age and older and young adults with relapsed or refractory, systemic anaplastic large cell lymphoma (ALCL) that is ALK-positive. • Ordered for treatment when the indication has been recognized by the National Comprehensive Cancer Network (NCCN) Drugs and Biologics Compendium with a category of Evidence and Consensus of 1, 2A, or 2B. <p><u>Non-Small Cell Lung Cancer (NSCLC):</u></p> <ul style="list-style-type: none"> • The cancer is either: metastatic, recurrent, or advanced. • The tumor is one of the following: <ul style="list-style-type: none"> ○ ALK-positive ○ ROS1-positive ○ Positive for mesenchymal-epithelial transition (MET) amplification ○ Positive for MET exon 14 skipping mutation 	<p>1. Patient does not show evidence of progressive disease while on Xalkori therapy.</p> <p>2. Approval Duration: 12 months.</p>

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p><u>Soft Tissue Sarcoma/Uterine Neoplasms:</u></p> <ul style="list-style-type: none"> • Diagnosis of inflammatory myofibroblastic tumor (IMT); and • Presence of ALK translocation. <p><u>Histiocytic Neoplasms:</u></p> <ul style="list-style-type: none"> • Diagnosis of one of the following: <ul style="list-style-type: none"> ○ Erdheim-Chester Disease ○ Langerhans Cell Histiocytosis ○ Rosai-Dorfman Disease • Used as targeted therapy ALK-fusion. <p><u>Central Nervous System (CNS) Cancers:</u></p> <ul style="list-style-type: none"> • Diagnosis of metastatic brain cancer from NSCLC; and • Tumor is ALK-positive OR ROS1-positive. <p><u>Anaplastic Large Cell Lymphoma:</u></p> <ul style="list-style-type: none"> • Tumor is ALK-positive; and • Disease is relapsed or refractory <p><u>Melanoma:</u></p> <ul style="list-style-type: none"> • Diagnosis of metastatic or unresectable cutaneous melanoma; and • Disease is ROS1 gene fusion-positive; and • Used as second-line or subsequent therapy for disease progression, intolerance, and/or projected risk of progression with BRAF-targeted therapy. <p>2. Medication ordered by an Oncologist. 3. Approval Duration: 12 months.</p>	
dabrafenib (Tafinlar) capsules 50mg, 75mg	1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • treatment of patients with unresectable or metastatic melanoma with BRAF V600E or V600K mutations as detected by an FDA-approved test 	<p><u>Limitations of use:</u></p> <ul style="list-style-type: none"> • Tafinlar is not indicated for treatments of patients with colorectal cancer because of

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> • adjuvant treatment of patients with melanoma with BRAF V600E or V600K mutations, as detected by an FDA-approved test, and involvement of lymph node(s), following complete resection. • treatment of patients with metastatic non-small cell lung cancer (NSCLC) with BRAF V600E mutation as detected by an FDA-approved test. • treatment of patients with locally advanced or metastatic anaplastic thyroid cancer (ATC) with BRAF V600E mutation and with no satisfactory locoregional treatment options. • Treatment of adult and pediatric patients ≥ 6 years of age with unresectable or metastatic solid tumors with BRAF V600E mutation who have progressed following prior treatment and have no satisfactory alternative treatment options. • Treatment of pediatric patients ≥ 1 year of age with low-grade glioma (LGG) with BRAF V600E mutation who require systemic therapy. <ol style="list-style-type: none"> 2. Medication ordered by an Oncologist 3. Approval duration: 12 months 	<p>known intrinsic resistance to BRAF inhibition.</p> <ul style="list-style-type: none"> • Tafinal is not indicated for treatment of patients with wild-type BRAF solid tumors • The indication for treatment of adult and pediatric patients 6 years of age and older with unresectable or metastatic solid tumors with BRAF V600E mutation continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trials.
dalfampridine (Ampyra) ER tablets 10mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • To Improve walking in adult patients with multiple sclerosis (MS). 2. Patient age ≥ 18 years. 3. Patient is currently receiving therapy with an agent to reduce progression of multiple sclerosis. 4. Patient does not have history of seizure. 5. Patient has appropriate renal function; CrCl > 50 ml/min. 6. Must be able to walk 25 feet within 8 to 45 seconds at baseline. 	<ol style="list-style-type: none"> 1. Improvement in walking speed as demonstrated by T25FW as compared with baseline. 2. Approval duration: 12 months.

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ol style="list-style-type: none"> 7. Must have a baseline gait assessment by PT within 90 days of beginning Ampyra. 8. Limited to 2 tablets per day. 9. Medication ordered by a Neurologist. 10. Initial approval for 3 months only after 3 months, must show improvement in walking speed must be documented to obtain continued approval. 	
daprodustat (Jesduvroq) tablets 1 mg, 2 mg, 4 mg, 6 mg, 8 mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Treatment of anemia that is caused by chronic kidney disease (CKD) in adults who have been on dialysis for at least 4 months. 2. Patient age \geq 18 years. 3. Patient on dialysis. 4. Pre-treatment hemoglobin level is $<$ 11 g/dL. 5. Serum transferrin saturation (TSAT) \geq 20% within prior 3 months. 6. Cannot use concomitantly with other erythropoiesis stimulating agents. 7. Maximum daily dose 24 mg per day. 8. Initial approval duration: 6 months. 	<ol style="list-style-type: none"> 1. Can not increase dose more frequently than once every 4 weeks. 2. Serum transferrin saturation (TSAT) \geq 20% within prior 3 months. 3. May not use concomitantly with other erythropoiesis stimulating agents. 4. After 24 weeks, if hemoglobin has not increased by \geq 1 g/dL, then therapy should be discontinued and cannot be approved. 5. Approval duration: 6 months.
darolutamide (Nubeqa) tablets 300mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • treatment of non-metastatic castration-resistant prostate cancer (mCRPC). • Metastatic hormone-sensitive prostate cancer (mHSPC) in combination with docetaxel. 2. Patient is \geq 18 years of age, AND 3. The medication is concurrently used with docetaxel OR the patient has completed docetaxel therapy. 4. The patient meets ONE of the following: <ul style="list-style-type: none"> • The medication is used concurrently with a gonadotropin-releasing hormone (GnRH) agonist, or 	<ol style="list-style-type: none"> 1. Patient has not shown disease progression. 2. Patient has not experienced unacceptable toxicity. 3. Patient should also receive a GnRH analog concurrently OR have had a bilateral orchiectomy. 4. Treatment may continue even if a cycle of docetaxel is delayed, interrupted, or discontinued. 5. Approval Duration: 12 months.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> • The medication is used concurrently with degarelix SQ injection; or • Patient has bilateral orchiectomy. 5. Medication ordered by an Oncologist or Urologist. 6. Approval Duration: 12 months.	
denosumab (Prolia) injection 60mg/ml	1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • treatment of postmenopausal women with osteoporosis at high risk for fracture. • treatment to increase bone mass in men with osteoporosis at high risk for fracture. • treatment of glucocorticoid-induced osteoporosis in men and women at high risk for fracture. • treatment to increase bone mass in men at high risk for fracture receiving androgen deprivation therapy for non-metastatic prostate cancer. • treatment to increase bone mass in women at high risk for fracture receiving adjuvant aromatase inhibitor therapy for breast cancer. 2. Patient age ≥ 18 years of age. 3. Tried and failed, had adverse reaction to, or contraindication to formulary preferred products (e.g., alendronate, calcitonin nasal spray). 4. Baseline calcium and vitamin D level results, with plan to correct any identified deficiencies before treatment initiation. 5. Baseline dental exam completed, and any preventative dentistry performed before treatment initiation. 7. Concomitant use of calcium and vitamin D supplement required. 8. For patients with advanced kidney disease (eGFR <30 mL/minute/1.73 m ²), including dialysis-dependent	1. All initial criteria met. 2. Approval Duration: 12 months. 3. NOTE: drug discontinuation conveys an increased risk of fractures and would require transition to alternative agent based on clinical guidance.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>patients: evaluation for presence of chronic kidney disease-mineral disorder (CKD-MBD) must be completed prior to denosumab initiation. Treatment with denosumab in these patients should be supervised by a health care provider with expertise in the diagnosis and management of CKD-MBD.</p> <p>9. Authorization duration: 12 months.</p>	
<p>dextromethorphan/quinidine (Nuedexta) tablets 10mg-20mg</p>	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Treatment of pseudobulbar affect (PBA) 2. Patient age \geq 18 years. 3. Patient has been diagnosed with ONE of the following: <ul style="list-style-type: none"> • Amyotrophic lateral sclerosis (ALS) • Alzheimer’s disease • Multiple sclerosis (MS) • Parkinson’s disease • Stroke • Traumatic brain injury 4. The baseline Center for Neurologic Study-Lability Scale (CNS-LS) score must be $>$ 13. 5. Dose must not exceed 2 capsules per day. 6. Prescribed by or in consultation with a neurologist. 7. Initial Authorization period is limited to 6 months. 	<ol style="list-style-type: none"> 1. Documentation of positive clinical response to therapy. 2. Authorization period is up to 12 months. <p>Limitations of Use: The following indications are considered experimental and cannot be approved:</p> <ul style="list-style-type: none"> • Heroin detoxification • Levodopa-induced Dyskinesia in Parkinson’s Disease • Neuropathic pain • Psychosis-Related Aggression • Treatment Resistant Depression
<p>diazepam nasal spray (Valtoco) doses: 5mg, 10mg, 15mg, 20mg doses</p>	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Treatment of active seizures (non-status epilepticus) in patients diagnosed with a seizure disorder. 2. Patient age \geq6 and $<$12 years of age (if 12 years or older, Nayzilam [midazolam] is preferred for seizure treatment as it is provided as a flat dose). 3. Requested dose is appropriate based on current weight (dosing below is for children 6-11 years): 	<ol style="list-style-type: none"> 1. Initial approval criteria continue to be met. 2. Current patient weight is provided. 3. Requested dose is appropriate based on current weight. <p>Approval duration: 12 months</p>

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria										
	<table border="1" data-bbox="657 250 1251 488"> <thead> <tr> <th>Weight (kg)</th> <th>Dose (mg)</th> </tr> </thead> <tbody> <tr> <td>10 to <19</td> <td>5 mg, as 1x 5mg device</td> </tr> <tr> <td>19 to <38</td> <td>10 mg, as 1x 10mg device</td> </tr> <tr> <td>38 to <56</td> <td>15 mg, as 2x 7.5mg devices</td> </tr> <tr> <td>56 to 74</td> <td>20 mg, as 2x10 mg devices</td> </tr> </tbody> </table> <p>Approval duration: 12 months</p>	Weight (kg)	Dose (mg)	10 to <19	5 mg, as 1x 5mg device	19 to <38	10 mg, as 1x 10mg device	38 to <56	15 mg, as 2x 7.5mg devices	56 to 74	20 mg, as 2x10 mg devices	
Weight (kg)	Dose (mg)											
10 to <19	5 mg, as 1x 5mg device											
19 to <38	10 mg, as 1x 10mg device											
38 to <56	15 mg, as 2x 7.5mg devices											
56 to 74	20 mg, as 2x10 mg devices											
dulaglutide (Trulicity) 0.75 mg, 1.5 mg, 3 mg, 4.5 mg	<ol style="list-style-type: none"> Ordered for an approved indication for use: <ul style="list-style-type: none"> As an adjunct to diet and exercise to improve glycemic control in adults or pediatric patients ≥ 10 years of age with type 2 diabetes mellitus. In adult patients with T2DM for risk reduction of major cardiovascular events (cardiovascular death, nonfatal myocardial infarction, nonfatal stroke) in adults with type 2 diabetes mellitus who have established cardiovascular disease or multiple cardiovascular risk factors. Patient is aged ≥ 10 years of age. Patient has Type 2 Diabetes Mellitus ***NOTE: Type 1 DM does NOT qualify for coverage*** A1c or CGM Time In Range% (TIR) report within past 3 months. A urine albumin-to-creatinine ratio (uACR) within the previous 12 months. <p><u>Treatment of Type 2 Diabetes without regard to CVD risk factors:</u> The patient has an A1c (hemoglobin A1c) of ≥ 7.5 (TIR $\leq 60\%$) OR</p> <p><u>Treatment of Type 2 Diabetes with CVD as defined below:</u></p>	<p>Cannot be approved for indication of weight management.</p> <ol style="list-style-type: none"> Chart notes with A1c or CGM report with TIR% within previous 3 months. A urine albumin-to-creatinine ratio (uACR) within the previous 12 months. Documented positive clinical response defined as one of the following: <u>Baseline (pre-GLP-1) A1c was ≥ 8.0 and:</u> 4. A1c has decreased by $\geq 1\%$ since onset of therapy or TIR% was $\leq 55\%$ and has increased $\geq 10\%$ OR <u>Baseline (pre-GLP-1) A1c was ≥ 6.5 but < 8.0 and:</u> <ul style="list-style-type: none"> A1c or TIR% has improved Not eligible for renewal if A1c is unchanged or has increased or TIR% has decreased. Patient has not had medical intervention for <ul style="list-style-type: none"> Pancreatitis; or Severe gastrointestinal 										

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> • Pre-treatment A1c is ≥ 6.5 (TIR $\leq 70\%$) AND • BMI ≥ 27 kg/m² (documentation within previous 90 days of current height and weight) AND <p>Documentation submitted to show that the patient has at least ONE of the following:</p> <ul style="list-style-type: none"> • History of myocardial infarction; OR • Prior Stroke (ischemic or hemorrhagic stroke); OR • Peripheral arterial disease as evidenced by: <ul style="list-style-type: none"> ○ Intermittent claudication with ankle-brachial index (ABI) less than 0.85 (at rest); OR ○ Peripheral arterial revascularization procedure; OR ○ Amputation due to atherosclerotic disease. <p>6. May not be concurrently using:</p> <ul style="list-style-type: none"> • Any other GLP1 or GLP1/GIP combination drug (e.g., Mounjaro, Ozempic, Rybelsus, Saxenda, Soliqua, Victoza [liraglutide], Xultrophy or Zepbound). • Any DPP4i (e.g., alogliptin, Januvia (sitagliptin), Tradjenta (Linagliptin), Onglyza (saxagliptin)). • Agents for severe constipation: metoclopramide, Amitiza (lubiprostone), Linzess (linaclotide), Motegrity (prucalopride) or Trulance (plecanatide). <p>7. Prescriber attests that medication is prescribed in accordance with prescribing information, including screening for any black box warnings and all contraindications.</p>	<p>events. (e.g., hospitalization or new start GI motility agent).</p> <p>5. May not be concurrently using:</p> <ul style="list-style-type: none"> • ANY other GLP1 or GLP1/GIP combination drug (e.g., Mounjaro, Ozempic, Rybelsus, Saxenda, Soliqua, Trulicity, Victoza (liraglutide), Xultrophy or Zepbound) • ANY DPP4i (e.g., alogliptin, Januvia (sitagliptin), Onglyza (saxagliptin), or Tradjenta (linagliptin)). • Agents for severe constipation: metoclopramide, Amitiza (lubiprostone), Linzess (linaclotide), Motegrity (prucalopride) or Trulance (plecanatide). <p>6. PBM claims data shows consistent adherence as shown by no instance of a drug-free interval greater than 2 months at which time the patient would need to satisfy the initial criteria.</p> <p>5. Approval Duration: up to 12 months</p>

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>8. May not be approved for patients with:</p> <ul style="list-style-type: none"> • Any personal or family history of medullary thyroid carcinoma (MTC) or multiple endocrine neoplasia syndrome type 2 (MEN 2). • A history of confirmed pancreatitis, and/or • Current pregnancy <p>9. Cannot be approved for indication of weight management.</p> <p>10. Trulicity 0.75 mg is a starter dose and is limited to one, 28-day supply and then must be dose escalated to 1.5 mg per week dose UNLESS A1c ≤ 7.0 or TIR ≥ 65%</p> <p>11. Approval Duration: up to 12 months</p>	
<p>dupilumab (Dupixent) subcutaneous injection</p> <p>Pen-injector – 200 mg/1.14 ml Pen-injector – 300 mg/2 ml Prefilled syringe – 200mg/1.14 ml Prefilled syringe – 300 mg/2 ml</p>	<p>1. Prescribed for an FDA-approved indication for use.</p> <p>2. The dosage and frequency requested are aligned with FDA and manufacturer guidelines for patient-specific parameters:</p> <ul style="list-style-type: none"> • Patient age • Patient weight • Indication for use <p>3. The criteria for Dupixent are indication specific. Please review criteria for the patient-specific diagnosis.</p> <p>Atopic Dermatitis:</p> <ul style="list-style-type: none"> • Diagnosis of moderate-to-severe chronic atopic dermatitis; AND • History of failure, contraindication, or intolerance to TWO of the following therapeutic classes of topical therapies (document drug, date of trial, and/or contraindication to medication). 	<p>Renewal criteria are indication specific. Please review criteria for the patient-specific diagnosis.</p> <p>Atopic Dermatitis:</p> <ul style="list-style-type: none"> • Documentation of a positive clinical response to therapy; AND • Patient is not Dupixent concurrent with either of the following: <ul style="list-style-type: none"> ○ Biologic immunomodulator (e.g., Adbry (tralokinumab-ldrm), Ebglyss (lebrikizumab), etc.) ○ Janus kinas inhibitor (e.g., Rinvoq (Upadacitinib), Xeljanz/XR (tofacitinib), Opzelura (topical ruxolitinib), Cibinqo (abrocitinib)); AND • Prescribed by a Dermatologist,

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> ○ Medium-high, or very-high potency topical corticosteroid (e.g. mometasone, fluocinolone acetonide, fluocinonide). ○ Topical calcineurin inhibitor (e.g., tacrolimus or pimecrolimus). ○ Phosphodiesterase-4 Enzyme Inhibitor, e.g. Zoryve (roflumilast), Eucrisa (crisaborole); AND ● Patient is not receiving Dupixent concurrent with either of the following: <ul style="list-style-type: none"> ○ Biologic immunomodulator (e.g., tralokinumab-ldrm); ○ Janus kinas inhibitor (e.g., Ebglyss (lebrikizumab), Rinvoq (Upadacitinib), Xeljanz/XR (tofacitinib), Opzelura (topical ruxolitinib), Cibinco (abrocitinib) ● UNLESS Patient has $\geq 25\%$ skin involvement and topical management is not feasible. ● Prescribed by or a Dermatologist, Allergist or Immunologist. ● Approval Duration: 12 months. <p>Asthma, moderate to severe eosinophilic:</p> <ul style="list-style-type: none"> ● Diagnosis of moderate-to-severe asthma; AND ● Classification of asthma as uncontrolled or inadequately controlled as defined by at least ONE of the following: <ul style="list-style-type: none"> ○ Poor symptom control (e.g., Asthma Control Questionnaire [ACQ] score consistently > 1.5 or Asthma Control Test [ACT] score consistently < 20). 	<p>Allergist, or Immunologist.</p> <ul style="list-style-type: none"> ● Approval Duration: 12 months. <p>Asthma:</p> <ul style="list-style-type: none"> ● Documentation of positive clinical response as demonstrated by at least ONE of the following: <ul style="list-style-type: none"> ○ Reduction in frequency of exacerbations. ○ Decreased utilization of rescue medications. ○ Increased in % predicted FEV1 from pre-treatment baseline. ○ Reduction in severity or frequency of asthma-related symptoms (e.g., wheezing, SOB, coughing) ○ Reduction in oral corticosteroid requirements; AND ○ Dupixent is being used in combination with an ICS-containing maintenance medication (e.g. fluticasone/salmeterol, Breo Ellipta, budesonide/formoterol, Trelegy); AND ○ Patient is not receiving Dupixent in combination with any of the following: Anti-interleukin-5 therapy

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> ○ Two or more bursts of systemic corticosteroids for at least 3 days each in the previous 12 months ○ Asthma-related emergency treatment (e.g., ER visit, hospital admission, or unscheduled physicians' office visit for nebulizer or other urgent treatment) ○ Airflow limitation (e.g., after appropriate bronchodilator withhold forced expiratory volume in 1 second [FEV1] < 80% predicted) ○ Patient is currently dependent on oral corticosteroids for the treatment of asthma; AND ● ONE of the following: <ul style="list-style-type: none"> ○ Submission of medical records documenting that asthma is an eosinophilic phenotype as defined by a baseline (pre-dupilumab treatment) peripheral blood eosinophil level ≥ 150 cells/μL; OR ○ Patient is currently dependent on oral corticosteroids for the treatment of asthma; AND ○ Dupixent will be used in combination with ONE of the following: <ul style="list-style-type: none"> ▪ On maximally dosed combination inhaled ICS/LABA inhaler (e.g., Advair, AirDuo, Symbicort, Breo, etc); OR ○ Combination therapy including BOTH of the following: 	<p>(e.g. mepolizumab, reslizumab, benralizumab); Anti-IgE therapy (e.g. omalizumab); and/or Thymic stromal lymphopoietin (TSLP) inhibitor (e.g. Tezepelumab); AND</p> <ul style="list-style-type: none"> ● Prescribed by an Allergist, Immunologist, or Pulmonologist. ● Approval Duration: 12 months. <p><u>Chronic Obstructive Pulmonary Disease (COPD)</u></p> <ul style="list-style-type: none"> ● Documentation of positive clinical response to therapy as defined by at least one of the following criteria: <ul style="list-style-type: none"> ○ A reduction in moderate exacerbations (i.e., those requiring systemic steroids and/or antibiotics). ○ A reduction of severe exacerbations (i.e. those requiring hospitalization and requiring more than one day of observation in an emergency department or urgent care facility). ○ An improvement in baseline lung function as assessed by pre-bronchodilator

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> ▪ One maximally dosed ICS product (e.g. Alvesco, Asmanex, Qvar, etc); AND ▪ One additional asthma controller medication (e.g., LABA, montelukast or theophylline); AND • Patient is not receiving Dupixent in combination with ANY of the following: <ul style="list-style-type: none"> ○ Anti-interleukin-5 therapy (e.g. Nucala, Cinqair, Fasentra). ○ Anti-IgE therapy (e.g., Xolair). ○ Thymic stromal lymphopoietin (TSLP) inhibitor (e.g. Tezspire); AND • Prescribed by or a Dermatologist, Allergist, Immunologist or Pulmonologist. • Approval Duration: 12 months. <p><u>Chronic Obstructive Pulmonary Disease (COPD)</u></p> <ul style="list-style-type: none"> • May be approved as add-on therapy in refractory disease who are inadequately controlled on standard therapies. • Patient age ≥ 18 years. • Diagnosis of COPD confirmed by spirometry (FEV1/FVC < 0.7). • Patient is actively using a triple therapy inhaler (e.g. Breztri or Trelegy). Active use is confirmed by pharmacy claims data showing ≥ 65% of utilization over time in the previous 6 months. • Patient has had 2 or more moderate exacerbations (i.e. symptoms requiring treatment with systemic glucocorticosteoids) OR at least 1 hospitalization for COPD exacerbation in previous 12 months, AND 	<p>forced expiratory volume (FEV1).</p> <ul style="list-style-type: none"> • Approval Duration: 12 months. <p><u>Chronic Rhinosinusitis with Nasal Polyposis</u></p> <ul style="list-style-type: none"> • Documentation of positive clinical response to Dupixent therapy; AND • Patient will continue to receive Dupixent as add-on maintenance therapy in combination with intranasal corticosteroids; AND • Patient is not receiving Dupixent in combination with ANY of the following: Anti-interleukin-5 therapy (e.g. mepolizumab, reslizumab, benralizumab); Anti-IgE therapy (e.g. omalizumab); and/or Thymic stromal lymphopoietin (TSLP) inhibitor (e.g. Tezepelumab); AND • Prescribed by an Allergist, Immunologist, or Pulmonologist. • Approval Duration: 12 months. <p><u>Eosinophilic Esophagitis:</u></p> <ul style="list-style-type: none"> • Documentation of positive clinical response to Dupixent therapy as evidenced by improvement in at least ONE of the following from baseline: <ul style="list-style-type: none"> ○ Symptoms ○ Histologic measures

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> • Pre-treatment blood eosinophil count \geq 300 cells/microliter. • Prescribed by or in consultation with a Pulmonologist. • Approval Duration: 12 months. <p>Chronic Rhinosinusitis with Nasal Polyposis</p> <ul style="list-style-type: none"> • Diagnosis with chronic rhinosinusitis with nasal polyposis (CRSwNP) defined by ALL of the following: <ul style="list-style-type: none"> ○ TWO or more of the following symptoms for longer than a 12-week duration: <ul style="list-style-type: none"> ▪ Nasal mucopurulent discharge ▪ Nasal obstruction, blockage or congestion ▪ Facial pain, pressure and/or fullness ▪ Reduction or loss of sense of smell; AND ○ ONE of the following findings using nasal endoscopy and/or sinus computed tomography: <ul style="list-style-type: none"> ▪ Purulent mucus or edema in the middle meatus or ethmoid regions ▪ Polyps in the nasal cavity or the middle meatus ▪ Radiographic imaging demonstrating mucosal thickening or partial or complete opacification of paranasal sinuses; AND ○ ONE of the following: <ul style="list-style-type: none"> ▪ Presence of bilateral nasal polyposis 	<ul style="list-style-type: none"> ○ Endoscopic measures; AND • Patient is not receiving Dupixent in combination with ANY of the following: Anti-interleukin-5 therapy (e.g. mepolizumab, reslizumab, benralizumab); Anti-IgE therapy (e.g. omalizumab); and/or Thymic stromal lymphopoietin (TSLP) inhibitor (e.g. Tezepelumab); AND • Prescribed by a Gastroenterologist or Allergist. • Approval Duration: 6 months. <p>Prurigo Nodularis</p> <ul style="list-style-type: none"> • Documentation of positive clinical response to Dupixent therapy; AND • Patient is not receiving Dupixent in combination with EITHER of the following: <ul style="list-style-type: none"> ○ Biologic immunomodulator (e.g., Adbry) OR ○ Janus kinase inhibitor (e.g., Rinvoq, Xeljanz/XR, Opzelura, Cingiqo); AND • Prescribed by a Dermatologist, an Allergist, or an Immunologist. <p>Approval Duration: 12 months.</p>

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> <ul style="list-style-type: none"> ▪ Patient has previously required surgical removal of bilateral nasal polyps; AND ○ ONE of the following: <ul style="list-style-type: none"> ▪ Patient has required prior sinus surgery ▪ Patient has required systemic corticosteroids for CRSwNP in the previous 2 years ▪ Patient has been unable to obtain symptom relief after trial of TWO of the following classes of agents: <ul style="list-style-type: none"> ➤ Nasal saline irrigations ➤ Intranasal corticosteroids ➤ Antileukotriene agents; AND • Patient will receive Dupixent as add-on maintenance therapy in combination with intranasal corticosteroids; AND • Patient is NOT receiving Dupixent in combination with ANY of the following: <ul style="list-style-type: none"> ○ Anti-interleukin-5 therapy (e.g. mepolizumab, reslizumab, benralizumab); ○ Anti-IgE therapy (e.g. omalizumab); and/or ○ Thymic stromal lymphopoietin (TSLP) inhibitor (e.g. Tezepelumab); AND • Prescribed by an Allergist, an Immunologist, an Otolaryngologist, or a Pulmonologist. • Approval Duration: 12 months. Eosinophilic Esophagitis: <ul style="list-style-type: none"> • Diagnosis of Eosinophilic Esophagitis; AND • Patient aged ≥ 2 years of age; AND 	

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> • Patient is experiencing symptoms related to esophageal dysfunction (e.g., dysphagia, food impaction, chest pain that is centrally located and may not respond to antacids, gastroesophageal reflux disease-like symptoms/refractory heartburn, upper abdominal pain); AND • Submission of clinical documentation indicating eosinophil-predominant inflammation on esophageal biopsy, consisting of a peak value of ≥ 15 intraepithelial eosinophils per high-power field (HPF) or 60 eosinophils per mm^2; AND • Secondary causes of esophageal eosinophilia have been ruled out; AND • Mucosal eosinophilia is isolated to the esophagus and symptoms have persisted after an 8-week trial of at least ONE of the following: <ul style="list-style-type: none"> ○ Proton pump inhibitor ○ Topical (esophageal) corticosteroids (e.g., budesonide, fluticasone); AND • Patient is not receiving Dupixent in combination with any of the following: <ul style="list-style-type: none"> ○ Anti-interleukin-5 therapy (e.g. mepolizumab, reslizumab, benralizumab); ○ Anti-IgE therapy (e.g. omalizumab); and/or ○ Thymic stromal lymphopoietin (TSLP) inhibitor (e.g. Tezepelumab); AND • Prescribed by either a Gastroenterologist or Allergist. • Approval Duration: 6 months. <p>Prurigo Nodularis</p> <ul style="list-style-type: none"> • Diagnosis of prurigo nodularis; AND • Patient has ≥ 20 nodular lesions; AND 	

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> • History of failure, contraindication, or intolerance to previous prurigo nodularis treatment(s) (e.g., topical corticosteroids, topical calcineurin inhibitors, topical capsaicin); AND • Patient is not receiving Dupixent with EITHER of the following: <ul style="list-style-type: none"> ○ Biologic immunomodulator (e.g. Adbry); OR ○ Janus kinase inhibitor (e.g., Rinvoq, Xeljanz/XR, Opzelura, Cibinqo); AND • Prescribed by a Dermatologist, Allergist, or Immunologist. <p>2. Approval Duration: 6 months.</p>	
eculizumab (Soliris) injection – 300mg/30ml	<p>1. Ordered for an approved indication for use:</p> <ul style="list-style-type: none"> • treatment of adult and pediatric patients ≥ 1 month of age with paroxysmal nocturnal hemoglobinuria (PNH). • treatment of adult and pediatric patients ≥ 1 month of age with atypical hemolytic uremic syndrome (aHUS) to inhibit complement-mediated thrombotic microangiopathy (TMA). • treatment of adult patients with generalized myasthenia gravis (gMG) who are anti-acetylcholine receptor (AChR+) antibody positive. • treatment of neuromyelitis optica spectrum disorder (NMOSD). <p>2. FDA approved patient age.</p> <p>3. Documentation to support diagnosis:</p> <p>PNH:</p> <ul style="list-style-type: none"> • Flow cytometric confirmation of PNH type III red cells; AND 	<p>1. Clinical documentation must be provided to confirm that current criteria are met and that the medication is providing clinical benefit.</p> <p>2. PNH:</p> <ul style="list-style-type: none"> • Age ≥ 18 years • Decrease in serum LDH from pre-treatment baseline. • NO dual therapy with another PA medication for PNH (e.g., Empaveli or Ultomiris). <p>aHUS:</p> <ul style="list-style-type: none"> • Decrease in serum LDH from pre-treatment baseline. • Patient does not have Shiga toxin E.coli related hemolytic uremic syndrome (STEC-HUS).

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> ○ Patient had at least one transfusion in the preceding 24 months; OR ○ Documented history of major adverse thrombotic vascular events from thromboembolism; OR ○ Patient has high disease activity defined as lactic dehydrogenase (LDH) level ≥ 1.5 times the upper limit of normal with one of the following symptoms: weakness, fatigue, hemoglobinuria, abdominal pain, dyspnea, hemoglobin, 10 g/dL, a major vascular event, dysphagia, or erectile dysfunction. <ul style="list-style-type: none"> ● History of failure to/contraindication or intolerance to Empaveli therapy; ● Patient age < 18 years or currently pregnant. <p>aHUS:</p> <ul style="list-style-type: none"> ● Common causes of aHUS have been ruled out, including infectious causes of HUS and thrombotic thrombocytopenic purpura (TTP). ● Ultomiris is not indicated for the treatment of patients with Shiga toxin E. coli related hemolytic uremic syndrome (STEC-HUS). ● Must present with the following symptoms: <ul style="list-style-type: none"> ○ Hemoglobin < 10 g/dL ○ Platelets, 150,000/mm³ ○ Documented evidence of hemolysis, such as elevated LDH levels, decreased haptoglobin level or schistocytosis. ○ Increased serum creatinine OR currently undergoing dialysis. <p>gMG:</p>	<ul style="list-style-type: none"> ● NO dual therapy with another PA medication for aHUS (e.g., Ultomiris). <p>gMG:</p> <ul style="list-style-type: none"> ● Age ≥ 18 years ● Improvement and maintenance of at least a 2-point improvement (reduction) in the MG-ADL score from pre-treatment baseline and reduction of signs and symptoms of MG required to show clinical benefit. ● NOTE: dose escalation of immunosuppressive therapy, or additional rescue therapy from baseline to treat MG or exacerbation of symptoms during use is considered treatment failure. ● Not receiving in combination with Empaveli or Ultomiris. <p>NMOSD:</p> <ul style="list-style-type: none"> ● Documentation to demonstrate positive clinical response from baseline as demonstrated by both of the following: ● Reduction in the number and/or severity of relapses or signs and symptoms of NMOSD; and ● Maintenance, reduction or discontinuation of dose(s) of any

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> • Patient has not failed previous course of Soliris or Ultomiris therapy; • Positive serologic test for anti-AChR antibodies; AND • One of the following: <ul style="list-style-type: none"> ○ History of abnormal neuromuscular transmission test demonstrated by single-fiber electromyography (SFEMG) or repetitive nerve stimulation OR ○ History of positive anticholinesterase test (e.g. edrophonium chloride test) OR ○ Pt has demonstrated improvement in MG signs on oral cholinesterase inhibitors as assessed by the treating neurologist; AND • Patient has MGFA clinical classification of II, III, or IV at initiation of treatment; AND • Patient has Myasthenia gravis-specific activities of daily living scale (MG-ADL) total score ≥ 6 at initiation of treatment; AND • One of the following: <ul style="list-style-type: none"> ○ History of failure to at least two immunosuppressive agents over the previous 12-months (e.g., azathioprine, mtx, cyclosporine, mycophenolate); OR ○ History of failure to at least one immunosuppressive therapy and has required four or more courses of plasmapheresis/plasma exchanges, and/or intravenous immune globulin over the previous 12 months without symptom control <p>NMOSD:</p>	<p>baseline immunosuppressive therapy prior to starting Soliris.</p> <ul style="list-style-type: none"> • NOTE: dose escalation of immunosuppressive therapy, or additional rescue therapy from baseline to treat MG or exacerbation of symptoms during use is considered treatment failure. • Not receiving Soliris therapy in combination with any of the following: Disease modifying therapies for the treatment of MS (e.g., Gilenya, Tecfidera, Ocrevus, etc.); Anti-IL6 therapy (e.g., Actemra, Enspryng), B-cell depletion therapy (e.g., rituximab, Uplizna). <p>3. Approval Duration: 3 months.</p>

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> • Documentation to support diagnosis of NMOSD by a neurologist confirming: <ul style="list-style-type: none"> ○ Optic neuritis; or ○ Acute myelitis; or ○ Area postrema syndrome: episode of otherwise unexplained hiccups or nausea and vomiting; or ○ Acute brainstem syndrome; or ○ Symptomatic cerebral syndrome with NMOSD-typical brain lesions; AND • Positive serologic test for anti-aquaporin-4 immunoglobulin G (AQP4-IgG)/NMO-IgG antibodies; and • Diagnosis of multiple sclerosis or other diagnoses have been ruled out; and • Patient has not failed a previous course of Soliris therapy; and • History of failure of, contraindication, or intolerance to rituximab therapy; and • History of at least two relapses in the previous 12-months; or • History of at least three relapses during the previous 24-months, at least one relapse within the previous 12-months; and • Not receiving Soliris therapy in combination with any of the following: Disease modifying therapies for the treatment of MS (e.g., Gilenya, Tecfidera, Ocrevus, etc.); Anti-IL6 therapy (e.g., Actemra, Enspryng), B-cell depletion therapy (e.g., rituximab, Uplizna). <p>4. Recommended vaccinations at least 2 weeks prior to administration of first dose Soliris.</p>	

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	5. Cannot be used in combination with other medications in the same class, such as Ultomiris. 6. Medication ordered by Hematologist, Nephrologist, or Oncologist. 7. Approval Duration: 3 months	
elacestrant (Orserdu)	1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Treatment of postmenopausal women or adult men, with estrogen receptor (ER)-positive, human epidermal growth factor receptor 2 (HER2)-negative, <i>ESR1</i>-mutated advanced or metastatic breast cancer with disease progression following at least one line of endocrine therapy. 2. For patient aged < 19 years of age: <ul style="list-style-type: none"> • If criteria in #1 are met, approval shall be granted for 12 months. For patients ≥ 19 years of age: <ul style="list-style-type: none"> • Diagnosis of breast cancer that is either advanced or metastatic. • The cancer is ER+, HER2(-) and has a confirmed <i>ESR1</i> gene mutation. • Patient is either male or a postmenopausal female. • Disease has progressed following at least one line of endocrine therapy. 3. Authorization Duration: 12 months.	1. Patient does not show evidence of progressive disease while on Orserdu therapy. 2. Approval Duration: 12 months.
elagolix, estradiol, and norethindrone acetate (Oriahnn) capsules 300mg/1mg/0.5mg	1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • the management of heavy menstrual bleeding associated with uterine leiomyomas (fibroids) in premenopausal women. 2. The patient is biologically female and premenopausal. 3. Negative pregnancy test result.	Renewal Criteria: 1. Patient continues to meet initial approval clinical criteria. 2. Approval Duration: up to 12 months, to allow the patient to reach the maximum lifetime duration of 24 months.

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ol style="list-style-type: none"> 4. Must have tried and failed or have a contraindication to using: <ul style="list-style-type: none"> • Combined estrogen/progestin-containing contraceptives (oral pills, transdermal patch, vaginal ring); OR • Levonorgestrel-releasing intrauterine device (IUD; e.g. Kyleena, Liletta, Mirena, Skyla); OR • Progestin-only oral contraceptive pills; OR • Tranexamic acid capsules. 5. Confirmation the patient does not have any contraindication to use Oriahnn: history of thrombotic/thrombolic disorders and women at increased risk of these events, including women > 35 years who smoke and women with uncontrolled hypertension. 6. The patient has not exceeded 24 cumulative months of treatment with an elagolix-containing medicine (e.g. Oriahnn, Orilissa) or a relugolix-containing medicine (e.g. Myfembree). 7. The patient is not concurrently using another gonadotropin-releasing hormone (GnRH) antagonist medicine (Orilissa, Myfembree) 8. Approval Duration: <ul style="list-style-type: none"> • 12 months; OR • The remaining duration to equal 24 months of cumulative therapy as listed in criteria #6, if less than 12 months remain. 	<p>Limitations of use: Due to risk of bone density loss that may not be reversible following discontinuation, limit duration of treatment to 24 months.</p>
elexacaftor, ivacaftor, and tezacaftor (Trikafta) tablets Therapy Pack	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • treatment of cystic fibrosis (CF) in patients \geq 2 years with at least one F508del mutation in the CFTR gene or a mutation in the CFTR gene that is responsive based on in vitro data. 	<ol style="list-style-type: none"> 1. Provider attestation of continued benefit without adverse drug effects. 2. Provider attestation of continued monitoring as appropriate. 3. Approval Duration: 12 months.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ol style="list-style-type: none"> 2. If the patient’s genotype is unknown, an FDA-cleared CF mutation test should be used to confirm the presence of at least one F508del mutation or a mutation that is responsive based on in vitro data. 3. Patient age \geq 2 years. 4. Provider attestation of baseline and subsequent evaluation and monitoring as appropriate and indicated by the FDA-approved product labeling (provider must submit documentation). 5. Provider justification of necessity of medication change if currently stable on another CF regimen and asymptomatic. 6. Medication ordered by a Pulmonologist. 7. Approval duration: 12 months. 	
<p>eltrombopag (Promacta) 12.5 mg, 25 mg packets for oral suspension; 12.5 mg, 25 mg, 50 mg, 75 mg tablets</p> <p>**note that Alvaiz (eltrombopag choline) is non-formulary</p>	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Treatment of thrombocytopenia in patients aged 1 year and older with persistent or chronic immune (idiopathic) thrombocytopenia (ITP) who have had an insufficient response to corticosteroids, immunoglobulins, or splenectomy. • Treatment of thrombocytopenia in patients with chronic Hepatitis C to allow the initiation and maintenance of interferon-based therapy. • First-line treatment of severe aplastic anemia in patients \geq 2 years of age in combination with standard immunosuppressive therapy. • Treatment of patients with severe aplastic anemia who have had an insufficient response to immunosuppressive therapy. 2. Pre-treatment platelet count lab results. 3. May not be treated concurrently with other thrombopoietin receptor agonists (e.g. Alvaiz, 	<p>May not be treated concurrently with other thrombopoietin receptor agonists (e.g. Alvaiz, Doptelet, Nplate, Mulpleta) or with spleen tyrosine kinase inhibitors (e.g. Tavalisse).</p> <p>Chronic Immune Thrombocytopenia (ITP):</p> <ol style="list-style-type: none"> 1. Documentation of positive clinical response to Promacta. 2. Current platelet count. 3. Approval Duration: 3 months for patients with current platelet counts less than $50 \times 10^9/L$ for whom the platelet count is not sufficient to prevent clinically important bleeding and who have not received a maximal dose of Promacta for at least 4 weeks. 4. Approval Duration: 12 months for patients: <ul style="list-style-type: none"> • with current platelet counts less

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>Doptelet, Nplate, Mulpleta) or with spleen tyrosine kinase inhibitors (e.g. Tavalisse).</p> <p>4. Promacta should be used only in patients with ITP whose degree of thrombocytopenia and clinical condition increase the risk for bleeding and should not be used in an attempt to normalize platelet counts.</p> <p>Chronic Immune Thrombocytopenia (ITP):</p> <ol style="list-style-type: none"> 1. Diagnosis of ITP; AND 2. Patient has had an insufficient response to a previous treatment (e.g. corticosteroids, immunoglobulins, thrombopoietin receptor agonists, splenectomy). 3. Pre-treatment platelet counts. 4. Untransfused platelet count at any point prior to the initiation of Promacta less than $30 \times 10^9/L$ or $30 \times 10^9/L$ to $50 \times 10^9/L$ with symptomatic bleeding (e.g. significant mucous membrane bleeding, GI bleeding or trauma) or risk factors for bleeding. 5. Ordered by or in consultation with a hematologist or oncologist. 6. Approval Duration: 6 months. <p>Chronic Hepatitis C-associated Thrombocytopenia:</p> <ol style="list-style-type: none"> 1. Diagnosis of chronic hepatitis C-associated thrombocytopenia; AND 2. One of the following: <ul style="list-style-type: none"> • Planning to initiate and maintain interferon-based treatment, OR • Currently receiving interferon-based treatment. • Ordered by or in consultation with a provider specializing in infectious disease, gastroenterology, hepatology, or transplant. • Approval Duration: 6 months. <p>Aplastic Anemia:</p> <ol style="list-style-type: none"> 1. Diagnosis of severe aplastic anemia; AND 2. Patient aged ≥ 2 years; AND 	<p>than $50 \times 10^9/L$ for whom the platelet count is sufficient to prevent clinically important bleeding OR</p> <ul style="list-style-type: none"> • for current platelet counts of $50 \times 10^9/L$ to $200 \times 10^9/L$, OR • patients with current platelet count $> 200 \times 10^9/L$ to $\leq 400 \times 10^9/L$ for whom dosing Promacta will be adjusted to achieve a platelet count sufficient to avoid clinically important bleeding. <p>Chronic Hepatitis C-associated Thrombocytopenia:</p> <ol style="list-style-type: none"> 1. Documentation of positive clinical response to Promacta; AND 2. Patient is currently on antiviral interferon therapy for treatment of chronic hepatitis C. 3. Approval Duration: 6 months <p>Aplastic Anemia:</p> <ol style="list-style-type: none"> 1. Documentation of positive clinical response to Promacta. 2. Current platelet counts. 3. Approval Durations: <ul style="list-style-type: none"> • Up to 4 months total for patients with current platelet counts less than $50 \times 10^9/L$ who have not received appropriately titrated therapy with Promacta for at least 16 weeks. • Up to 4 months total for patients with current platelet counts $< 50 \times 10^9/L$ who are transfusion-

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	3. ONE of the following: <ul style="list-style-type: none"> Used in combination with standard immunosuppressive therapy (e.g., Atgam [antithymocyte globulin equine], Thymoglobulin [antithymocyte globulin rabbit], cyclosporine), OR History of failure, contraindication, or intolerance to at least one course of immunosuppressive therapy (e.g. cyclosporine, Atgam, Thymoglobulin). 4. Approval Duration: 6 months.	independent. <ul style="list-style-type: none"> 12 months for patient with current platelet counts of $50 \times 10^9/L$ to $200 \times 10^9/L$. 12 months for patients with current platelet count $> 200 \times 10^9/L$ to $\leq 400 \times 10^9/L$ for whom dosing Promacta when dose adjusted to achieve and maintain the appropriate target platelet count.
enzalutamide (Xtandi) tablets 80mg	1. Ordered for an approved indication for use: <ul style="list-style-type: none"> castration-resistant prostate cancer for patients > 18 years of age. Metastatic, castration-sensitive prostate cancer (mCRPC) for patients > 18 years of age. 2. Patient is ≥ 18 years of age, AND 3. The patient meets ONE of the following: <ul style="list-style-type: none"> The medication is used concurrently with a gonadotropin-releasing hormone (GnRH) agonist, or Patient has bilateral orchiectomy; or Patient has non-metastatic, castration-sensitive cancer and a biochemical recurrence and at high-risk for metastasis (PSA doubling time ≤ 9 months). 4. Medication ordered by an Oncologist or Urologist. 5. Approval Duration: 12 months.	Patients receiving Xtandi should also receive a gonadotropin-releasing hormone (GnRH) analog concurrently or should have had bilateral orchiectomy. Examples of GnRH agonists include: leuprolide acetate, Lupron Depot (leuprolide acetate intramuscular injection), Trelstar (triptorelin pamoate intramuscular injection), Zoladex (goserelin acetate subcutaneous implant), Vantas (histrelin acetate subcutaneous implant).
etanercept (Enbrel; Enbrel Mini; Enbrel Sureclick) injection 25mg/0.5mL, 50mg/mL syringes 50mg/mL autojector	1. Ordered for an approved indication for use and following the indication-specific criteria listed below: Rheumatoid Arthritis (RA): <ol style="list-style-type: none"> Diagnosis of moderately to severely active RA. Patient meets one of the following: 	Renewal Criteria applies to all approved indications described in the initial criteria column: 5. Documentation of positive clinical response to therapy.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> • Has history of failure to a 3-month trial of one non-biologic disease modifying anti-rheumatic drug (DMARD) (e.g. methotrexate, leflunomide, sulfasalazine, hydroxychloroquine) at maximally indicated doses, unless contraindicated or clinically significant adverse effects are experienced (document drug, date, and duration of trial). OR • Patient has been previously treated with a targeted immunomodulator FDA-approved for the treatment of RA as documented by claims history or submission of medical records. (document drug, date and duration of therapy). (e.g. Cimzia, adalimumab, Simponi, Olumiant, Rinvoq, Xeljanz), OR <ol style="list-style-type: none"> 2. Patient is not receiving Enbrel in combination with another targeted immunomodulator (as listed in #2). 3. Prescribed by or in consultation with a rheumatologist. 4. Approval Duration: 12 months. <p><u>Polyarticular Juvenile Idiopathic Arthritis (PJIA)</u></p> <ol style="list-style-type: none"> 1. Diagnosis of moderately to severely active PJIA. 2. Patient is not receiving Enbrel concurrently with another targeted immunomodulator. 3. Prescribed by or in consultation with a rheumatologist. 4. Approval Duration: 12 months <p><u>Psoriatic Arthritis (PsA)</u></p> <ol style="list-style-type: none"> 1. Diagnosis of Psoriatic Arthritis. 2. The patient meets ONE of the following: <ul style="list-style-type: none"> • Patient has history of failure to a 3-month trial of methotrexate a maximally indicated dose, unless 	<ol style="list-style-type: none"> 6. Patient is not receiving Enbrel concurrently with another targeted immunomodulator (e.g. Cimzia, adalimumab, Simponi, Olumiant, Rinvoq, Xeljanz). 7. Approval Duration: 12 months.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>contraindicated or clinically significant adverse effects are experienced (with documentation of trial dates and details), OR</p> <ul style="list-style-type: none"> • Patient has been previously treated with a targeted immunomodulator that is FDA-approved for the treatment of PsA as documented in claims history or submission of medical records that include the name of the drug, dates, and duration of therapy, OR <p>3. Prescribed by or in consultation with a rheumatologist or dermatologist.</p> <p>4. Approval Duration: 12 months.</p> <p><u>Plaque Psoriasis:</u></p> <ol style="list-style-type: none"> 1. Diagnosis of moderate to severe plaque psoriasis. 2. Patient has greater than or equal to 3% body surface area involvement, palmoplantar, facial, genital involvement or severe scalp psoriasis, AND <ul style="list-style-type: none"> • History of failure to one of the following topical therapies unless contraindicated or clinically significant adverse effects are experienced with documentation included: <ul style="list-style-type: none"> ○ Topical corticosteroids ○ Vitamin D analogs (calcitriol, calcipotriene) ○ Tazarotene ○ Calcineurin inhibitors (tacrolimus/pimecrolimus) ○ Anthralin ○ Coal tar; AND • History of failure to a 3-month trial of methotrexate at maximally indicated dose unless contraindicated or clinically adverse effects occurred, OR 	

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	3. Patient has been previously treated with a targeted immunomodulator FDA-approved for the treatment of plaque psoriasis as documented by claims history or submission of medical records that include the name of the drugs, dates and duration of therapy. 4. Prescribed by or in consultation with a dermatologist. 5. Approval Duration: 12 months.	
etrasimod (Velsipity) tablets 2 mg	1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Treatment of ulcerative colitis (UC), in adults with moderately to severely active disease. 2. Patient is \geq 18 years of age. 3. Patient has had a trial of one systemic agent for ulcerative colitis. (e.g., 6-mercaptopurine, azathioprine, cyclosporine, tacrolimus, or a corticosteroid such as prednisone or methylprednisolone). Note: a trial of one biologic is considered a trial of systemic agent for ulcerative colitis. 4. Patient is not being treated concurrently with a biologic or targeted synthetic disease-modifying antirheumatic drug (DMARD) for UC. (e.g., adalimumab, infliximab, sarilumab, abatacept, rituximab, mirkizumab, ustekinumab, apremilast, ozanimod, or similar). 5. Medication is prescribed by or in consultation with a gastroenterologist. 6. Initial Approval Duration: 12 months.	1. Patient exhibits a positive clinical response by at least one objective measure from baseline. (e.g., fecal calprotectin levels, C-reactive protein, endoscopic assessment, and/or decreased utilization of corticosteroids OR 2. Patient has a documented clinical improvement in at least one subjective measure from baseline (e.g., decreased pain, fatigue, stool frequency, and/or rectal bleeding). 3. Approval duration: 12 months.
evinacumab-dgnb (Evkeeza) injection 345mg/2.3ml, 1200mg/8ml	1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • An adjunct to other low-density lipoprotein-cholesterol (LDL-C) lowering therapies for the treatment of adult and pediatric patients, \geq 12 	1. Meets all initial criteria 2. Must provide documentation of laboratory information to support continued use (full lipid panel) and

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>years of age, with homozygous familial hypercholesterolemia (HoFH).</p> <ol style="list-style-type: none"> 2. Documented genetic test confirming homozygous familial hypercholesterolemia (HoFH). 3. Baseline laboratory information required (full lipid panel, genetic testing, negative pregnancy test and documentation of use/counseling regarding contraception to prevent pregnancy) 4. Prior trial/failure and/or documented intolerance to one high potency statin (atorvastatin, rosuvastatin) and concurrent ezetimibe. <ul style="list-style-type: none"> • Must provide laboratory data to support failure/intolerance (full lipid panel, creatinine kinase). • If failure, but no intolerance, lipid lowering therapy should be continued with aa statin and/or ezetimibe. 5. Dosing 15 mg/kg IV every 4 weeks. 6. Initial Approval Duration: 6 months. 	<p>continued use of concurrent therapies to lower cholesterol</p> <ol style="list-style-type: none"> 3. Renewal Approval Duration: 3 months
<p>factor VIIa, recombinant human (NovoSeven RT) injection 1mg, 2mg, 5mg, 8mg</p>	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • treatment of bleeding episodes and perioperative management in adults and children with hemophilia A or B with inhibitors, congenital Factor VII (FVII) deficiency, and Glanzmann’s thrombasthenia with refractoriness to platelet transfusions, with or without antibodies to platelets. • treatment of bleeding episodes and perioperative management in adults with acquired hemophilia. 2. Diagnosis of congenital factor VIII deficiency confirmed by blood coagulation testing. 	<ol style="list-style-type: none"> 1. Patient continues to meet indication-specific criteria 2. Absence of unacceptable toxicity from drug; and 3. Any dose increases must be supported by an acceptable clinical rationale (i.e., weight gain, half-life study results, increase in breakthrough bleeding when patient is fully adherent to therapy, etc). 4. The cumulative amount of medication that the patient has on-hand will be considered. The authorization will

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ol style="list-style-type: none"> 3. Confirmation that patient has acquired inhibitors to Factor VIII 4. Used as treatment in at least one of the following: <ul style="list-style-type: none"> • Control and prevention of acute bleeding episodes; or • Perioperative management; or • Routine prophylaxis to prevent or reduce the frequency of bleeding episodes when the following criteria are also met: <ul style="list-style-type: none"> ○ Patient has at least two documented episodes of spontaneous bleeding into joints; or ○ Patient has documented trial and failure of Immune Tolerance Induction (ITI). 5. When ordered for Hemophilia B: <ul style="list-style-type: none"> • Diagnosis of congenital Factor IX deficiency has been confirmed by blood coagulation testing; and • Confirmation that patient has acquired inhibitors to Factor IX. 6. When ordered for Congenital Factor VII Deficiency: <ul style="list-style-type: none"> • Diagnosis confirmed by blood coagulation testing. 7. When ordered for Glanzmann’s Thromboasthnia: <ul style="list-style-type: none"> • Diagnosis confirmed by blood coagulation testing; and • The use of platelet transfusions is known or suspected to be ineffective or contraindicated. 8. Medication ordered by a Hematologist. 9. Approval Duration: <ul style="list-style-type: none"> • For perioperative management of bleeding: 1 month. • All other indications: up to 3 months. 	<p>allow up to 5 doses on hand for the treatment of acute bleeding episodes as needed for the duration of the authorization.</p> <ol style="list-style-type: none"> 5. Renewal duration: 3 months.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
factor VIII, recombinant human pegylated (Jivi) injection 500 unit, 1000unit, 2000unit, 3000unit	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • On-demand treatment and control of bleeding episodes in adults and adolescents ≥ 12 years of age with hemophilia A. • Perioperative management of bleeding. • Routine prophylaxis to reduce the frequency of bleeding episodes. 2. Patient has previously received Factor VIII replacement therapy. 3. Not for the treatment of von Willebrand disease. 4. Medication ordered by a Hematologist. 5. Authorization Duration: 12 months. 	<ol style="list-style-type: none"> 1. Documentation of positive clinical response to Jivi therapy. 2. Authorization Duration: 12 months.
fecal microbiota capsules, oral (Vowst)	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • To prevent recurrence of <i>Clostridioides difficile</i> infection (CDI) in individuals ≥ 18 years of age following antibacterial treatment for recurrent CDI. 2. Patient has had three or more episodes of CDI within previous 12 months (including most recent episode). 3. Patient has recent episode of recurrent CDI with all of the following: <ul style="list-style-type: none"> ○ At least 3 unformed stools per day for 2 consecutive days ○ Stool test confirming the presence of <i>C. difficile</i> toxin or toxigenic <i>C. difficile</i>. ○ An adequate clinical response (i.e., resolution of symptoms) following standard of care antibiotic therapy (e.g., vancomycin + metronidazole, fidaxomicin) 4. Patient does not have ANY of the following: <ul style="list-style-type: none"> ○ Known or suspected toxic megacolon and/or known small bowel ileus OR 	<ol style="list-style-type: none"> 1. Use is limited to two treatment courses per lifetime. 2. Patient must meet the initial criteria for use. <p>Limitations of Use: VOWST is not indicated to treat CDI.</p>

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> ○ Admitted to, or expected to be admitted to an ICU for medical reasons, OR ○ Absolute neutrophil count < 500 cells/mL³ ○ History of major GI surgery within 3 months before treatment start (not including appendectomy or cholecystectomy) OR ○ History of total colectomy or bariatric surgery that disrupted the GI lumen OR ○ History of active inflammatory bowel disease (e.g. ulcerative colitis, Crohn’s disease, microscopic colitis) with diarrhea believed to be cause by active inflammatory bowel disease in the past 3 months. ○ History of fecal microbiota transplantation (FMT) within 3 months <p>5. The patient will not be using the requested agent in combination with Rebyota or Zinplava for the requested indication.</p> <p>6. Provider attests that patient will follow the bowel preparation protocol outlined in the package insert.</p> <p>7. Patient will not be taking a concurrent antibiotic.</p> <p>8. Prescribed by or in consultation with an infectious disease specialist.</p> <p>9. Approval is limited to 12 capsules per dispense; maximum of 24 capsules lifetime.</p>	
fentanyl (Duragesic) transdermal patch 12mcg/hr, 25mcg/hr, 37.5mcg/hr, 50mcg/hr, 62.5mcg/hr, 75mcg/hr, 87.5mcg/hr, 100mcg/hr	<p>1. Ordered for an approved indication for use:</p> <ul style="list-style-type: none"> ● management of pain in opioid-tolerant patients, severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate. Patients considered opioid- tolerant are those taking, for one week or longer, at least 60 mg oral 	<ul style="list-style-type: none"> ● All long-acting opioids require prior authorization (PA). The PA request form can be access using the following links: <p>OPIOID PRIOR AUTH FORM-DC</p>

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>morphine per day, 25 mcg transdermal fentanyl per hour, 30 mg oral oxycodone per day, 8 mg oral hydromorphone per day, 25 mg oral oxymorphone per day, 60 mg oral hydrocodone per day, or an equianalgesic dose of another opioid.</p> <ol style="list-style-type: none"> 2. Fully completed opioid PA form submitted. 3. Submission of clinical documentation from last office visit, dated within 3 months of the request. 4. Maximum approval duration is 6 months but may be reduced based on any of the criteria as outlined in Pharmacy Policy 219.DC Opioid Prescription Prior Authorization. 	
fezolinetant (Veozah) tablets 45mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Treatment of moderate to severe vasomotor symptoms due to menopause. 2. Patient must be a perimenopausal or post-menopausal female 3. Documentation of baseline bloodwork to evaluate for hepatic function and injury including ALT, AST and serum bilirubin (total and direct) before initiation of treatment. 4. Provider attests to monitoring liver function tests at 3-months, 6-months, and 9-months after starting therapy. 5. Patient must not have cirrhosis. 6. Patient does not have severe renal impairment (GFR < 30 ml/min) or end-stage renal disease. 7. The medication must not be used concomitantly with CYP1A2 inhibitors (e.g., acyclovir, allopurinol, amiodarone, cimetidine, clarithromycin, duloxetine, famotidine, fluoroquinolones, fluvoxamine, mexiletine, oral contraceptives, verapamil, 	Renewal Criteria: <ol style="list-style-type: none"> 1. All criteria listed for initial approval AND: 2. Documented improvement of symptoms 3. Documentation of liver function tests monitoring during first year of treatment with labs within previous 3 months. 4. Renewal duration: 12 months

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	zafirlukast, zileuton). 8. Patient must have treatment failure, intolerance, or contraindication to at least one menopausal hormone therapy. 9. Initial approval period: 9 months	
finerenone (Kerendia) tablets 10mg, 20mg	1. Ordered for approved indication: <ul style="list-style-type: none"> • to reduce the risk of sustained eGFR decline, end stage kidney disease, cardiovascular death, non-fatal myocardial infarction, and hospitalization for heart failure in adult patients with chronic kidney disease (CKD) associated with type 2 diabetes (T2D). 2. PA SUBMISSION REQUIREMENTS: <ul style="list-style-type: none"> • Serum potassium \leq 5.0 mEq/L • eGFR \geq 25 mL/min/1.73 m² • Urine albumin-to-creatinine ratio \geq 30 mg/g • Concomitant use with maximum tolerated doses of ACE-Inhibitor or ARB unless intolerant to or contraindicated. 3. Failed trial or contraindication to at least one formulary SGLT2i. 4. Approval Duration: 3 months	1. All initial criteria for approval; AND 2. Dosing appropriate based on 4-week potassium laboratory check. <ul style="list-style-type: none"> • 20 mg daily if Potassium \leq 4.8 • 10 mg daily if K⁺ between 4.8-5.5 • Interrupt therapy if K⁺ > 5.5, may restart at 10 mg daily when potassium is \geq 5.0 3. Approval duration: 12 months
fosdenopterin (Nulibry) injection 9.5mg	1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • To reduce mortality risk in patients with molybdenum cofactor deficiency (MoCD) Type A. • Diagnosis confirmed by genetic testing. 2. Will not be used in combination with other substrate replacement therapy (e.g., recombinant cyclic pyranopterin monophosphate, etc.); AND 3. Must be prescribed by, or in consultation with, a specialist in medical genetics or pediatric neurology. 4. Diagnosis of MoCD Type A is confirmed by molecular	1. Patient continues to meet initial approval criteria as listed. 2. Absence of unacceptable toxicity from the drug (e.g., severe phototoxicity, clinically significant infection). 3. Disease response compared to pre-treatment baseline as evidenced by the following:

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>genetic testing, by a mutation in the <i>MOCS1</i> gene suggestive of disease.</p> <p>5. Patient has biochemical features suggestive of MoCD Type A (i.e., elevated sulfites in urine, low serum uric acid, elevated urinary xanthine and hypoxanthine) and will be treated presumptively while awaiting genetic confirmation; and</p> <p>6. Patient has baseline values for the following:</p> <ul style="list-style-type: none"> • Elevated urinary s-sulfocysteine (SSC) normalized to creatinine; and • Clinical notes regarding signs and symptoms of disease which may include, but are not limited to, seizure frequency/duration, growth, and developmental milestones. <p>7. Approval Duration: 3 months</p>	<ul style="list-style-type: none"> • Reduction in urinary SSC normalized to creatinine; and • Stabilization or improvement in one or more signs and symptoms of disease including, but not limited to, seizure frequency/duration, growth, achievement of developmental milestones; OR <p>4. Patient initiated therapy as an inpatient based on presumptive diagnosis of MoCD Type A which was subsequently confirmed by genetic testing; AND patient is responding to therapy compared to one or more pre-treatment baseline parameters which prompted the workup for MoCD.</p> <p>5. Approval Duration: 3 months</p>
fostamatinib disodium hexahydrate (Tavalisse) tablets 100mg, 150mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • the treatment of thrombocytopenia in adult patients with chronic immune thrombocytopenia (ITP) when a prior treatment for ITP has not worked well enough. 2. Patient age ≥ 18 years. 3. Patient is not on hemodialysis. 4. Max dose: 150 mg 2 times daily with goal platelets ≥ 50 x 10⁹/mmcp/L. 5. Medication ordered by a Hematologist. 6. Initial Approval Duration: 3 months. 	<ol style="list-style-type: none"> 1. Documentation of improved symptoms and attestation of lab parameters. 2. Renewal approval duration: 12 months

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
furosemide subcutaneous injection device (Furoscix) 80mg/10ml	<ol style="list-style-type: none"> 1. Ordered for the treatment of congestion due to fluid overload in adults with NYHA Class II/III chronic heart failure. 2. Patient has CrCl > 30 ml/min OR eGFR > 20 ml/min 3. Patient has been stable and is refractory to at least one of the following loop diuretics, at up to maximally indicated doses: <ul style="list-style-type: none"> ○ Furosemide oral tablets; 40-160 mg/day ○ Torsemide oral tablets; 50-100 mg/day ○ Bumetanide oral tablets; 3-10 mg/day 4. Documentation that member is a candidate for parenteral diuresis outside of the hospital, as defined by all of the following: <ul style="list-style-type: none"> ○ Oxygen saturation ≥ 90% on exertion ○ Respiratory rate < 24 breaths per minute ○ Resting heart rate < 100 beats per minute ○ Systolic blood pressure > 100 mmHg 5. Patient does not have anuria 6. Patient does not allergy to medical adhesives or furosemide. 7. Patient does not have hepatic cirrhosis or ascites. 8. Dose does not exceed 80 mg (1 cartridge) per day. 9. Prescribed by cardiologist 10. Limited to 8 kits every 30 days 11. Approval requires that patient is referred for MFC Case Management 12. Authorization Duration: 3 months 	Limitations of Use: <ul style="list-style-type: none"> • Furoscix is not indicated for emergency situations or in patients with acute pulmonary edema. • The On-Body Infusor will deliver only an 80-mg dose of Furoscix. • Patients must meet initial approval criteria for each request
gabapentin extended-release (Gralise) tablets 300mg, 600mg <small>*note, this is not the same as gabapentin enacarbil which is non-formulary.</small>	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • the management of Postherpetic Neuralgia (PHN). 	<ol style="list-style-type: none"> 1. Initial criteria continue to be met. 2. Approval duration: 12 months.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>Not interchangeable with other gabapentin products because of differing pharmacokinetic profiles that affect dosing frequency.</p> <ol style="list-style-type: none"> 2. Patient age \geq 18 years. 3. Patient CrCl > 30 ml/min; patient is not on hemodialysis. 4. Dose does not exceed 1800 mg per day. 5. Approval Duration: 12 months 	
gilteritinib (Xospata) tablets 40mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • the treatment of adult patients who have relapsed or refractory acute myeloid leukemia (AML) with a FLT3 mutation as detected by an FDA-approved test. 2. Medication ordered by an Oncologist 	<ul style="list-style-type: none"> •
glycopyrronium (Qbrexza) pad 2.4%	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • topical treatment of primary axillary hyperhidrosis in adults and pediatric patients \geq 9 years of age. 2. Patient age \geq 9 years. 3. Must have tried and failed OTC Clinical Strength antiperspirants and at least one prescription strength antiperspirant (ex: Drysol or Xerac AC) for at least 4 weeks and experienced inadequate efficacy 3. Documentation that symptoms are persistent despite previous treatment attempts and that the degree of symptomatology impacts quality of life must be clearly indicated in a recent (within past 6 months) clinical encounter note. 4. Qbrexza will only be applied to the axillae, and is not being used on other areas of the body. 5. Patient does not have any of the following conditions: <ul style="list-style-type: none"> • Glaucoma • Paralytic ileus 	<ol style="list-style-type: none"> 1. Patient has demonstrated benefit with Qbrexa. 2. Patient continues to meet initial approval criteria. 3. Approval Duration: 12 months

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> • Unstable cardiovascular status in acute hemorrhage • Severe ulcerative colitis • Toxic megacolon • Myasthenia gravis • Sjogren’s syndrome <p>6. Limited to 30 cloths per 30 days.</p> <p>7. Approval Duration: 12 months.</p>	
goserelin (Zoladex) implant 3.6mg, 10.8mg	<p>1. Ordered for an approved indication for use:</p> <ul style="list-style-type: none"> • palliative treatment of advanced carcinoma of the prostate. (3.6 mg and 10.8 mg) • in combination with flutamide for the management of locally confined stage T2b-T4 (Stage B2-C) carcinoma of the prostate. (3.6 mg and 10.8 mg) • management of endometriosis (3.6 mg) • palliative treatment of advanced breast cancer in pre- and peri-menopausal women. (3.6 mg) • to cause endometrial thinning agent prior to endometrial ablation for dysfunctional uterine bleeding. (3.6 mg) • management of endometriosis, including pain relief and reduction of endometriotic lesions for the duration of therapy. <p>Endometriosis:</p> <p>1. Contraindication, intolerance, or failure of initial treatment to BOTH of the following:</p> <ul style="list-style-type: none"> • Oral contraceptives or depot medroxyprogesterone; AND • Non-steroidal anti-inflammatory drugs; OR 	<p>Endometriosis:</p> <ul style="list-style-type: none"> • Can not be administered for more than 6 months lifetime maximum. <p>Endometrial thinning:</p> <ul style="list-style-type: none"> • Can not be administered for more than 6 months lifetime maximum. <p>Fertility Preservation:</p> <ul style="list-style-type: none"> • Patient currently receiving GnRH analog therapy for purpose of fertility preservation; and • Patient continues to receive a cytotoxic agent associated with primary ovarian insufficiency; and • Authorization duration: 12 months <p>Gender Affirming Care – Adolescents OR Gender Affirming Care – Transgender Adults:</p> <ul style="list-style-type: none"> • Approval Duration: 12 months.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> • Patient has had surgical ablation to prevent recurrence. <p>2. Approval Duration: Limited to 6 months.</p> <p><u>Endometrial Thinning/Dysfunctional Uterine Bleeding:</u></p> <ol style="list-style-type: none"> 1. For use prior to endometrial ablation; AND 2. Other causes of symptoms of bleeding are ruled out; AND 3. Patient has been prescribed the 3.6 mg implant; AND 4. Approval duration is for a maximum of 2 depots. <p><u>Fertility Preservation:</u></p> <p>Clinical studies do not support use for this indication, and cryopreservation is clinically preferred. Please attempt to redirect to cryopreservation. Only clinically appropriate as a potential adjunct to cryopreservation.</p> <p>May be medically necessary for treatment of fertility preservation when both of the following criteria are met:</p> <ul style="list-style-type: none"> • Patient is a pre-menopausal female. • Patient is receiving a cytotoxic agent associated with causing primary ovarian insufficiency, e.g., cyclophosphamide, procarbazine, vinblastine, cisplatin. • Approval Duration: 12 months. <p><u>Gender Affirming Care – Adolescents</u></p> <ol style="list-style-type: none"> 1. Prescribed by or in consultation with a medical provider experienced in transgender hormone therapy. 2. Approval Duration: 12 months <p><u>Gender Affirming Care – Transgender Adults</u></p> <ol style="list-style-type: none"> 1. Prescribed by or in consultation with a medical provider experienced in transgender hormone therapy. 	

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
ibrutinib (Imbruvica) capsules 140mg	<ol style="list-style-type: none"> 2. Approval Duration: 12 months 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Chronic lymphocytic leukemia (CLL) in adult patients who have received at least one prior therapy. • CLL in Adult patients with 17p deletion. • Waldenström’s macroglobulinemia in adult patients • Adult and pediatric patients ≥ 1 year of age with chronic graft versus host disease after failure of one or more lines of systemic therapy. 2. Medication ordered by an Oncologist. 3. Quantity limit: 4 tablets per day. 	<p>Limitations for use:</p> <ul style="list-style-type: none"> • Indications for Mantle Cell Lymphoma and Marginal Zone Lymphoma were voluntarily withdrawn, April 2023 <p><u>New dose modification guidelines adopted in December 2022:</u></p> <ol style="list-style-type: none"> 1. Therapy should be withheld for any new onset or worsening Grade 2 cardiac failure or Grade 3 cardiac arrhythmia. Once symptoms have resolved to Grade 1 cardiac failure or Grade 2 or lower cardiac arrhythmia, Imbruvica can be restarted at recommended adjusted doses.
icatibant acetate (Firazyr) injection 30mg/3ml	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • treatment of acute attacks of hereditary angioedema (HAE) in adults ≥ 18 years of age. 2. Patient age ≥ 18 years. 3. Prescribed for the treatment of acute HAE attacks. 4. Member has a C1 inhibitor deficiency or dysfunction as confirmed by laboratory testing and meets one of the following: <ol style="list-style-type: none"> 3. C1 inhibitor (C1-INH) antigenic level below the lower limit of normal as defined by the laboratory performing the test; OR <ul style="list-style-type: none"> • Normal C1-INH antigenic level and a low C1-INH functional level (functional C1-INH less than 50% or C1-INH functional level below the lower limit of normal as defined by the laboratory performing the test). 5. If not the criteria in #4 above, the patient has normal C1 inhibitor as confirmed by laboratory testing and 	<ol style="list-style-type: none"> 1. Patient meets initial approval criteria. 2. Submission of chart notes showing that Patient has experienced a reduction in severity and/or duration of attacks. 3. Prophylaxis should be considered based on the frequency and severity of attacks, comorbid conditions, and patient’s quality of life. 4. Approval Duration: 6 months.

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>meets one of the following criteria:</p> <ul style="list-style-type: none"> • Patient has an F12, angiotensin-converting enzyme, plasminogen, kininogen-1 (KNG1), heparan sulfate-glucosaminase 3-O-sulfotransferase 6 (HS3ST6), or myoferlin (MYOF) gene mutation as confirmed by genetic testing; or • Patient has a documented family history of angioedema and the angioedema was refractory to a trial of high-dose antihistamine therapy (i.e. cetirizine at 40 mg per day or the equivalent) for at least 30 days. <p>6. Not used in combination with other products indicated for the acute treatment of HAE attacks (e.g. Berinert, Kalbitor, or Ruconest).</p> <p>7. Medication ordered by an Allergist or ENT.</p> <p>8. Approval Duration: 6 months.</p>	
icosapent ethyl (E-EPA) capsules (Vascepa) 0.5gm, 1gm	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • As an adjunct to maximally tolerated statin therapy to reduce the risk of myocardial infarction, stroke, coronary revascularization, and unstable angina requiring hospitalization in adult patients with elevated triglyceride (TG) levels (≥ 150 mg/dL) AND <ul style="list-style-type: none"> ○ Established cardiovascular disease OR ○ Diabetes mellitus and 2 or more additional risk factors for cardiovascular disease • As an adjunct to diet to reduce TG levels in adult patients with severe (≥ 500 mg/dL) hypertriglyceridemia. 2. Age ≥ 45 years 3. Diagnosis of hypertriglyceridemia (pre-treatment TG level ≥ 150 mg/dl) AND 	Renewal Criteria: <ul style="list-style-type: none"> • Used for cardiovascular risk reduction • Documentation of positive clinical response to therapy • Patient is receiving maximally tolerated statin therapy. • Approval duration: 12 months

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>4. Patient is considered high or very high risk for cardiovascular disease (CVD) as evidenced by one of the following:</p> <ul style="list-style-type: none"> ○ Acute coronary syndrome ○ History of myocardial infarction ○ Stable or Unstable angina ○ Coronary or other arterial revascularization ○ Stroke ○ Transient ischemic attack ○ Peripheral arterial disease <p>5. OR, if not the criteria in #4:</p> <ul style="list-style-type: none"> ○ Type 2 Diabetes diagnosis AND TWO of the following: ○ Men ≥ 55 years and women ≥ 65 years ○ Cigarette smoker or stopped within past 3 months ○ Hypertension diagnosis ○ HDL-C ≥ 40 mg/dL for men or ≥ 50 mg/dL for women ○ High-sensitivity C-reactive protein > 3.0 mg/L ○ Creatinine clearance > 30 and < 60 mL/min ○ Retinopathy ○ Micro- or macro-albuminuria ○ Ankle-brachial index (ABI), 0.9 without symptoms of intermittent claudication <p>6. Patient has received at least 12 consecutive weeks of high-intensity statin therapy (Atorvastatin 40-80 mg; rosuvastatin 20-40 mg) OR BOTH OF THE FOLLOWING:</p> <ul style="list-style-type: none"> ● Intolerance to high-intensity statin as evidenced by ≥ 2 weeks of myalgia and/or 	

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>myositis AND</p> <ul style="list-style-type: none"> at least 12 consecutive weeks of low/moderate intensity statin therapy <p>7. Patient has been receiving at least 12 consecutive weeks of ezetimibe (Zetia) therapy as adjunct to maximally tolerated statin therapy, or contraindication or intolerance to ezetimibe OR has LDL-C less than 100 mg/dL while on maximally tolerated statin therapy.</p> <p>8. Approval duration: 12 months.</p>	
<p>idecabtagene vicleucel (Abecma) injection</p>	<ol style="list-style-type: none"> Ordered for an approved indication for use: <ul style="list-style-type: none"> To treat relapsed or refractory multiple myeloma in adults after ≥4 prior therapies, including an immunomodulatory agent, a proteasome inhibitor, and an anti-CD38 monoclonal antibody. Lymphodepleting chemotherapy (with fludarabine and cyclophosphamide) is ordered for administration for 3 days followed by Abecma dose infusion 2 days after completion of lymphodepleting therapy. Diagnosis of relapsed or refractory multiple myeloma (MM) Age ≥ 18 years Must have received at least 4 prior MM therapies (induction with or without hematopoietic stem cell transplant with or without maintenance therapy is considered a single regimen) Must have received an immunomodulatory drug (iMiD), proteasome inhibitor (PI), and an anti-CD38 antibody ECOG performance status of 0 or 1 HBV, HCV, and HIV screening within previous 30 days. 	<p>Limitations of use:</p> <ul style="list-style-type: none"> Will be approved for ONE treatment dose. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	9. Provider attestation: Drug specific baseline evaluation and monitoring completed according to package insert (CBC/CMP, screening for HBV, hepatitis C, HIV), patient is not pregnant and is using effective contraception, counseling/assessment of recent live vaccine use. 10. Monitor immunoglobulin levels, blood counts, and for cytokine release syndrome during and after therapy. 11. Medication ordered by Hematologist or Oncologist enrolled in ABECMA REMS and compliance with REMS program criteria.	
immune globulin subcutaneous (human) (Cutaquig) solution 1gm/6ml, 1.65gm/10ml, 2gm/12ml, 3.3gm/20ml, 4gm/24ml, 8gm/48ml	1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Replacement therapy for primary humoral immunodeficiency (PI) in adults and pediatric patients ≥ 2 years of age. • Prevention of bacterial infection in patients with hypogammaglobulinemia and/or recurrent bacterial infections with malignancy (e.g., B-cell chronic lymphocytic leukemia) or primary humoral immunodeficiency disorders. 2. Medication ordered by an Immunologist.	
interferon gamma-1b (Actimmune) injection 2 million IU/0.5ml	1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • To reduce frequency and severity of serious infections associated with chronic granulomatous disease (CGD). • To delay time to disease progression in patients with severe, malignant osteopetrosis (SMO). 2. Patient age is less than 19 years. 3. When prescribed for: <ul style="list-style-type: none"> • Chronic Granulomatous Disease (CGD); • Osteopetrosis; or 	1. Patient does not show evidence of progressive disease while on Actimmune. 2. Reauthorization is for 3 months. 3. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> • Primary Cutaneous Lymphomas when the patient has a diagnosis of: <ul style="list-style-type: none"> ○ Mycosis fungoides (MF) or ○ Sezary Syndrome (SS) <p>4. Approval Duration: 3 months.</p>	
ivacaftor (Kalydeco) tablets 150mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Treatment of cystic fibrosis (CF) in patients \geq 4 months who have one mutation in the CFTR gene that is responsive to ivacaftor potentiation based on clinical and/or <i>in vitro</i> assay data. 2. If the patient's genotype is unknown, an FDA-cleared CF mutation test should be used to detect the presence of a CFTR mutation followed by verification with bi-directional sequencing when recommended by the mutation test instructions for use. 3. Patient is not homozygous in the CFTR gene. 4. Patient age \geq 4 months. 5. Provider attestation of baseline and subsequent evaluation and monitoring as appropriate and as indicated in the FDA-approved labeling (provider must submit documentation). 6. Provider justification of necessity of medication change if currently stable on another CF regimen and asymptomatic. 7. Medication ordered by Pulmonologist. 8. Approval Duration: 12 months. 	<ol style="list-style-type: none"> 1. Provider attestation of continued benefit without adverse drug effects. 2. Provider attestation of continued monitoring as appropriate. 3. Approval Duration: 12 months.
ivermectin (Stromectol) tablets 3mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Strongyloidiasis of the intestinal tract (i.e., nondisseminated) strongyloidiasis due to the nematode parasite <i>Strongyloides stercoralis</i>. • Onchocerciasis due to the nematode parasite <i>Onchocerca volvulus</i>. 	<p>Limitations for use:</p> <ul style="list-style-type: none"> • At this time, outpatient use for COVID-19 treatment is prohibited. • Ivermectin has no activity against adult <i>Onchocerca volvulus</i> parasites.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	2. Cannot be used for outpatient COVID-19 treatment.	<ul style="list-style-type: none"> Ivermectin is not active against <i>L. loa</i> (adult worms).
larotrectinib (Vittrakvi) capsules 25mg, 100mg	<ol style="list-style-type: none"> Ordered for an approved indication for use: <ul style="list-style-type: none"> Treatment of adult and pediatric patients with solid tumors that have a neurotrophic receptor tyrosine kinase (NTRK) gene fusion without a known acquired resistance mutation, are metastatic or where surgical resection is likely to result in severe morbidity, and no satisfactory alternative treatments or that have progressed following treatment. The patient is being treated for one of the following solid tumors: soft tissue sarcoma, salivary gland, infantile fibrosarcoma, thyroid, lung, or gastrointestinal stromal tumors; and The tumor is positive for neurotrophic receptor tyrosine kinase (NTRK) gene fusion; AND The tumor is metastatic OR surgical resection of tumor will likely result in severe morbidity. Medication ordered by an Oncologist. Approval Duration: 6 months for first authorization. 	<ol style="list-style-type: none"> Patient continues to meet initial criteria. Patient has documented positive response to therapy as defined by stabilization of disease or decrease in tumor size or tumor spread. Absence of unacceptable toxicity from the drug (e.g. severe neurotoxicity, hepatotoxicity etc.) Approval Duration: 12 months
lebrikizumab (Ebglyss) injection 250mg/2mL syringe, autoinjector	<ol style="list-style-type: none"> Ordered for an approved indication: <ul style="list-style-type: none"> Treatment of patients ≥ 12 years of age who weigh at least 40 kg with moderate-to-severe atopic dermatitis whose disease is not adequately controlled with topical therapies or when those therapies are not advisable. Diagnosis of moderate-to-severe chronic atopic dermatitis; AND Patient age is ≥ 12 years Patient weight is ≥ 40 kg. 	<ol style="list-style-type: none"> Documentation of a positive clinical response to therapy; AND Patient is not using Ebglyss concurrent with any of the following: <ul style="list-style-type: none"> Biologic immunomodulator (e.g., Dupixent (dupilumab), Adbry (tralokinumab-ldrm); and/or Janus kinase inhibitor (e.g., Rinvoq (Upadacitinib), Xeljanz/XR (tofacitinib), Opzelura (ruxolitinib), Cibinqo (abrocitinib); AND

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ol style="list-style-type: none"> 5. History of failure, contraindication, or intolerance to TWO of the following therapeutic classes of topical therapies (document drug, dates of trial, and/or contraindication to medication). <ul style="list-style-type: none"> • Medium-high, or very-high potency topical corticosteroid (e.g. mometasone, fluocinolone acetonide, fluocinonide). • Topical calcineurin inhibitor (e.g. tacrolimus or pimecrolimus) • Phosphodiesterase-4 Enzyme Inhibitor (e.g. Zoryve (roflumilast), Eucrisa (crisaborole)). AND 6. Patient is not receiving Ebglyss concurrent with any of the following: <ul style="list-style-type: none"> • Biologic immunomodulators (e.g. Adbrey (tralokinumab-ldrm), or Dupixent). • Janus kinase inhibitors (e.g. Rinvoq (Upadacitinib), Xeljanz/XR (tofacitinib), Opzelura (ruxolitinib), Cibinqo (abrocitinib)). 7. Prescribed by or in consultation with a Dermatologist, Allergist, or Immunologist. 8. Approval Duration: 6 months. 	<ol style="list-style-type: none"> 3. Prescribed by or in consultation with a Dermatologist, Allergist, or Immunologist. 4. Approval Duration: 12 months.
lecanemab-irmb (Leqembi) intravenous solution 200 mg/2 ml, 500mg/5ml	<ol style="list-style-type: none"> 1. Ordered for an approved indication: <ul style="list-style-type: none"> • Treatment of Alzheimer disease; to be initiated in patients with mild cognitive impairment or mild dementia stage of disease, with confirmed presence of amyloid beta pathology prior to initiation of treatment. 2. Patient has signed informed consent on file. 3. Patient meets criteria for mild cognitive impairment (MCI) or mild AD dementia. 4. Patient has had an MRI scan within last 12 months. 5. Amyloid PET imaging and/or CSF analysis consistent 	Renewal Criteria: <ol style="list-style-type: none"> 1. Patient continues to meet criteria for initial approval. 2. Absence of unacceptable toxicity from drug AND 3. Patient has responded to therapy compared to pretreatment as evidenced by improvement, stability, or slowing in cognitive and/or functional impairment in one or more of the following (not all-inclusive):

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>with AD.</p> <ol style="list-style-type: none"> 6. Functional Assessment Staging Test Stage score of 2 to 4. 7. Mini-Mental State Examination score greater than 21, or St. Louis University Mental Status (SLUMS) score or Montreal Cognitive Assessment (MoCA) score of greater than 16. 8. Patient does not have any of the following risk factors for intracerebral hemorrhage: <ul style="list-style-type: none"> • prior cerebral hemorrhage greater than 1 cm in greatest diameter, more than 4 microhemorrhages, superficial siderosis, • evidence of vasogenic edema, • evidence of cerebral contusion, • aneurysm, • vascular malformation, • infective lesions, • multiple lacunar infarcts or stroke involving a major vascular territory, • and severe small vessel or white matter disease. 9. Ordered by a Board-certified neurologist, geriatric psychiatrist, or geriatrician who specializes in treating dementia. 	<p>ADAS-Cog 13; ADCS-ADL-MCI; MMSE: CDR-SB etc, AND</p> <ol style="list-style-type: none"> 4. Patient has not progressed to moderate or severe AD; AND 5. Patient has received a pre-5th, 7th, AND 14th infusion MRI for monitoring of Amyloid Related Imaging Abnormalities-edema (ARIA-E) and Amyloid Related Imaging Abnormalities hemosiderin (ARIA-H) microhemorrhages.
<p>leuprolide injection</p> <p>leuprolide acetate kit 1mg/0.2ml</p> <p>Eligard SQ injection 45 mg</p> <p>Lupron Depot IM injection 1-month (3.75mg, 7.5mg) 3-month (11.25mg, 22.5mg)</p>	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • palliative treatment of advanced carcinoma of the prostate. • in combination with flutamide for the management of locally confined stage T2b-T4 (Stage B2-C) carcinoma of the prostate. • management of endometriosis 	<p><u>Endometriosis:</u></p> <ol style="list-style-type: none"> 1. Can not be administered for more than 6 months lifetime maximum. <p><u>Fertility Preservation:</u></p> <ul style="list-style-type: none"> • Patient currently receiving GnRH analog therapy for purpose of fertility preservation; and • Patient continues to receive a cytotoxic agent associated with primary ovarian insufficiency; and

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
<p>4-month (30mg)</p> <p>Lupron Depot-PED IM injection kit</p> <p>1-month (7.5mg, 11.25mg 15mg)</p> <p>3-month (11.25mg, 30mg)</p> <p>6-month (45mg)</p>	<ul style="list-style-type: none"> • palliative treatment of advanced breast cancer in pre- and peri-menopausal women. • to cause endometrial thinning agent prior to endometrial ablation for dysfunctional uterine bleeding. • management of endometriosis, including pain relief and reduction of endometriotic lesions for the duration of therapy. <p><u>Endometriosis:</u></p> <p>1. Contraindication, intolerance, or failure of initial treatment to BOTH of the following:</p> <ul style="list-style-type: none"> • Oral contraceptives or depot medroxyprogesterone; AND • Non-steroidal anti-inflammatory drugs; OR • Patient has had surgical ablation to prevent recurrence. <p>2. Approval Duration: Limited to 6 months.</p> <p><u>Fertility Preservation:</u></p> <p>May be medically necessary for treatment of fertility preservation when both of the following criteria are met:</p> <ul style="list-style-type: none"> • Patient is a pre-menopausal female. • Patient is receiving a cytotoxic agent associated with causing primary ovarian insufficiency, e.g., cyclophosphamide, procarbazine, vinblastine, cisplatin. • Approval Duration: 12 months. <p><u>Gender Affirming Care – Adolescents</u></p> <p>1. Prescribed by or in consultation with a medical provider experienced in transgender hormone therapy.</p> <p>2. Approval Duration: 12 months</p> <p><u>Gender Affirming Care – Transgender Adults</u></p>	<ul style="list-style-type: none"> • Authorization duration: 12 months <p><u>Gender Affirming Care – Adolescents</u> OR <u>Gender Affirming Care – Transgender Adults:</u></p> <ul style="list-style-type: none"> • Approval Duration: 12 months. <p><u>Oncology Indications:</u></p> <ul style="list-style-type: none"> • Patient has positive clinical response and absence of unacceptable toxicity <p><u>Uterine Leiomyomata (Fibroids) –</u></p> <ul style="list-style-type: none"> • Treatment beyond 6 months requires combination with add-back therapy (progestin, estrogen + progestin, NSAID), and should only be considered if surgery is contraindicated, or in post-surgical patients with persistent pain inadequately controlled with empiric therapies (NSAIDs, OCPs)

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ol style="list-style-type: none"> 1. Prescribed by or in consultation with a medical provider experienced in transgender hormone therapy. 2. Approval Duration: 12 months <p><u>Oncology Indications:</u></p> <ol style="list-style-type: none"> 1. Prescribed by a hematologist/oncologist AND 2. The requested use is supported by the National Comprehensive Cancer Network (NCCN) clinical practice guidelines with a recommendation category level of 1 or 2A. 3. <u>Oncology Approval duration:</u> <ul style="list-style-type: none"> • Prostate cancer: up to 90 mg per 12 months. • Breast/ovarian cancer: up to 22.5 mg per 6 months; approval duration is up to 6 months. <p><u>Uterine Leiomyomata (Fibroids) –</u></p> <ol style="list-style-type: none"> 1. Lupron Depot formulation prescribed 2. Prescribed for use prior to surgery to reduce the size of fibroids to facilitate surgical procedure; OR 3. For the treatment of uterine leiomyomata-related anemia; AND 4. inadequate respond to iron therapy of one month duration; AND 5. For use prior to surgery 6. Approval Duration: 6 months total. 	
lifitegrast ophthalmic (Xiidra) drops 5% STEP THERAPY	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • the treatment of the signs and symptoms of dry eye disease (DED). 2. Must have tried and failed artificial tears AND cyclosporine (ophth) emulsion 0.05% (generic of Restasis). 3. Approval Duration: 12 months. 	

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
liraglutide (Victoza) injection 1.2 mg/day 2-pack pens (6 ml) 1.8 mg/day 3-pack pens (9 ml)	<ol style="list-style-type: none"> Ordered for an approved indication for use: <ul style="list-style-type: none"> An adjunct to diet and exercise to improve glycemic control in adults and pediatric patients aged 10 years and older with type 2 diabetes To reduce the risk of major adverse cardiovascular events in adults with type 2 diabetes and established cardiovascular disease. Patient has Type 2 Diabetes Mellitus ***NOTE: Type 1 DM does NOT qualify for coverage*** A1c or CGM Time in Range% (TIR) report within past 3 months. A urine albumin-to-creatinine ratio (uACR) within the previous 12 months. <p><u>Treatment of Type 2 Diabetes without regard to CVD risk factors:</u></p> <p>The patient has an A1c (hemoglobin A1c of ≥ 7.5 (TIR $\leq 60\%$))</p> <p style="text-align: center;">OR</p> <p>Treatment of Type 2 Diabetes with CVD as defined below:</p> <ul style="list-style-type: none"> Pre-treatment A1c is ≥ 6.5 (TIR $\leq 70\%$) AND BMI ≥ 27 kg/m² (documentation within previous 90 days current height and weight); AND <p>Documentation submitted to show that the patient has at least one of the following:</p> <ul style="list-style-type: none"> History of myocardial infarction; or Prior stroke (ischemic or hemorrhagic); or Symptomatic peripheral arterial disease (PAD) as evidenced by: <ul style="list-style-type: none"> Intermittent claudication with ankle-brachial index (ABI) less than 0.85 (at rest); OR Peripheral arterial revascularization procedure; OR 	<p>Cannot be approved for the indication of weight management.</p> <ol style="list-style-type: none"> Chart notes with A1c or CGM report with TIR% within previous 3 months. A urine albumin-to-creatinine ratio (uACR) within the previous 12 months. Documented positive clinical response defined as one of the following: <p><u>Baseline (pre-GLP1) A1c was ≥ 8.0 and:</u></p> <ul style="list-style-type: none"> A1c has decreased by $\geq 1\%$ since onset of therapy or TIR% was $\leq 55\%$ and has increased $\geq 10\%$ or A1c is ≤ 7.0 at initiation dose. <p><u>Baseline (pre-GLP1) A1c was ≥ 6.5 but < 7.5 and:</u></p> <ul style="list-style-type: none"> A1c or TIR% has improved. NOT eligible for renewal if A1c has increased or TIR% has decreased. May not be concurrently using: <ul style="list-style-type: none"> ANY other GLP1 or GLP1/GIP combination drug (e.g., Ozempic, Rybelsus, Saxenda, Soliqua, Trulicity, Victoza, Xultrophy or Zepbound) AND/OR ANY DPP4i (e.g., alogliptin, Januvia (sitagliptin), Onglyza (saxagliptin), or Tradjenta (linagliptin)). Agents for severe constipation: metoclopramide, Amitiza (lubiprostone), Linzess (linaclotide), Motegrity (prucalopride) or Trulance

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> ○ Amputation due to atherosclerotic disease. <p>6. May not be concurrently using:</p> <ul style="list-style-type: none"> • ANY other GLP1 or GLP1/GIP combination drug (e.g., Mounjaro, Ozempic, Rybelsus, Saxenda, Soliqua, Trulicity, Wegovy, Xultrophy or Zepbound). • ANY DPP4i (e.g., alogliptin, Januvia (sitagliptin), Tradjenta (Linagliptin), Onglyza (saxagliptin)). • Agents for severe constipation: metoclopramide, Amitiza (lubiprostone), Linzess (linaclotide), Motegrity (prucalopride) or Trulance (plecanatide). <p>7. Prescriber attests that medication is prescribed in accordance with prescribing information, including screening for any black box warnings and all contraindications.</p> <p>8. May not be approved for patients with:</p> <ul style="list-style-type: none"> • Personal or family history of medullary thyroid carcinoma (MTC) or in patients with Multiple Endocrine Neoplasia syndrome type 2 (MEN 2). • Current pregnancy; and/or • A history of confirmed pancreatitis. <p>9. Cannot be approved for indication of weight management.</p> <p>10. Dose escalation in accordance with manufacturer guidelines required. Initial dose is 0.6 mg once daily for 1 week, then must increase to 1.2 mg daily as the 0.6 mg dose does not provide effective glycemic control.</p> <p>11. Quantity Limits:</p> <ul style="list-style-type: none"> • 1.2 mg daily dose is limited to one-pack containing 2 pens (6 ml) per 30 days. 	<p>(plecanatide).</p> <p>6. PBM claims data shows consistent adherence as shown by no instance of a drug-free interval greater than 2 months at which time the patient would need to satisfy the initial criteria.</p> <p>7. Approval Duration: 12 months</p>

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> 1.8 mg daily dose is limited to one pack containing 3 pens (9 ml) per 30 days. 12. Maximum Approval Duration: 12 months.	
lisdexamfetamine (Vyvanse) capsules 10mg, 20mg, 30mg, 40mg, 50mg, 60mg, 70mg chewables 10mg, 20mg, 30mg, 40mg, 50mg, 60mg STEP THERAPY	1. Ordered for an approved indication for use: <ul style="list-style-type: none"> Attention Deficit Hyperactivity Disorder (ADHD) in children ≥ 6 years of age. Moderate to Severe Binge Eating Disorder (BED) in adults. 3. Step therapy: <ul style="list-style-type: none"> <u>ADHD</u>: at least 4-week trial of an amphetamine salt combination AND a 4-week trial of methylphenidate. <u>BED</u>: at least 12-week trial of a serotonin reuptake inhibitor (SSRI) 	
lisocabtagene maraleucel (Breyanzi) injection 70,000,000 cells	1. Ordered for an approved indication for use: <ul style="list-style-type: none"> Treatment of adult patients with large B-cell lymphoma (LBCL) including diffuse large B-cell lymphoma (DLBCL) not otherwise specified (including DLBCL arising from indolent lymphoma), high-grade B-cell lymphoma, primary mediastinal large B-cell lymphoma, and follicular lymphoma grade 3B, who have: <ol style="list-style-type: none"> refractory disease to first-line chemoimmunotherapy or relapse within 12 months of first-line chemoimmunotherapy; or refractory disease to first-line chemoimmunotherapy or relapse after first-line chemoimmunotherapy and are not eligible for hematopoietic stem cell transplantation (HSCT) due to comorbidities or age; or 	<p style="text-align: center;">Not applicable. Maximum approval, one treatment course per lifetime.</p> <p>Limitations of Use:</p> <ul style="list-style-type: none"> BREYANZI is not indicated for the treatment of patients with primary central nervous system lymphoma.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ol style="list-style-type: none"> c. relapsed or refractory disease after two or more lines of systemic therapy. 2. Age ≥ 18 years of age. 3. Prescriber attestation that all baseline evaluations have been done, and no contraindications to use are present. 4. Prescriber attests that subsequent appropriate evaluation and monitoring will be done based on the package insert. 5. Dose: 50-110 x 10⁶ CAR positive viable T cells, one time dose. 6. Medication ordered by an Oncologist or Hematologist. 7. Approval limited to once per lifetime. 	
lomitapide (Juxtapid) capsules 5mg, 10mg, 20mg, 30mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • An adjunct to a low-fat diet and other lipid-lowering treatments, including LDL apheresis where available, to reduce LDL-C, total cholesterol, apolipoprotein B, and non-HDL-C in patients with homozygous familial hypercholesterolemia. 2. Patient age ≥ 18 years. 3. Documentation of baseline LFTs (including ALT, AST, alkaline phosphatase and total bilirubin) prior to initiation of treatment. 4. Prescriber attestation that a low-fat diet (<20% of energy from fat) has been initiated. 5. Prior trial, failure, insufficient response, and/or documented intolerance to preferred lipid lowering treatments including statin + ezetimibe, or Praluent. 6. Medication ordered by a REMS registered cardiologist or endocrinologist. 	<ol style="list-style-type: none"> 1. Meets all initial approval criteria. 2. Attestation of continued benefit without significant adverse drug effects. 3. Laboratory data (full lipid panel) submitted to support continued use. 4. Renewal Duration: 12 months

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
loncastuximab tesirine-lpyl (Zynlonta) solution 10mg	7. Approval Duration: 12 months. 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> the treatment of adult patients with relapsed or refractory large B-cell lymphoma after two or more lines of systemic therapy, including diffuse large B-cell lymphoma (DLBCL) not otherwise specified, DLBCL arising from low-grade lymphoma, and high-grade B-cell lymphoma. 2. Patient is at least 18 years of age. 3. Used as a single-agent therapy. 4. Patient has not received prior anti-CD19 therapy (e.g., tafasitamab, CAR-T) or patient previously received anti-CD19 therapy and re-biopsy indicates CD-19 positive disease; and 5. Patient does not have graft-versus-host disease; and 6. Patient has not had an autologous stem cell transplant (ASCT) within 30 days or allogeneic stem cell transplant within 60 days prior to start of therapy; and 7. Medication ordered by an Oncologist. 8. Approval Duration: 6 months	1. Patient continues to meet initial criteria for use. 2. Positive disease response from treatment defined as stabilization of disease or decrease in size of tumor or tumor spread. 3. Absence of unacceptable drug toxicity. 4. Approval Duration: 6 months.
lotilaner 0.25% solution (Xdemvy) 2.5mg/ml	1. Ordered for an approved indication for use: <ul style="list-style-type: none"> Treatment of Demodex blepharitis in adults 2. Patient aged ≥ 18 years of age. 3. Diagnosis of Demodex blepharitis, AND 4. Patient demonstrates at least one of the clinical signs of Demodex infestation: <ul style="list-style-type: none"> Cylindrical cuff at the root of the eyelashes (collarettes), lid margin erythema, eyelash anomalies (misdirected lashes); AND 	<ul style="list-style-type: none"> At this time, there is no clinical evidence to show benefit beyond 6 weeks of treatment.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ol style="list-style-type: none"> 5. Patient demonstrates two of the following symptoms of Demodex blepharitis in at least one eye: <ul style="list-style-type: none"> • Itching/burning, • foreign body sensation, • crusting/matter lashers • blurry vision • discomfort/irritation; AND 6. Clinical documentation indicates the patient has been educated about good eyelid hygiene practices. 7. Patient has not undergone more than 1 6-week treatment in the previous 12 months. 8. Written by or in consultation with an ophthalmologist or optometrist. 9. Approval limited to 1 bottle (10 ml) per 12 months. 	
lumacaftor/ivacaftor (Orkambi) tablets 100mg-125mg, 200mg-125mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • the treatment of cystic fibrosis (CF) in patients aged 1 year and older who are homozygous for the F508del mutation in the CFTR gene. 2. If the patient’s genotype is unknown, an FDA-cleared CF mutation test should be used to confirm the presence of the F508del mutation on both alleles of the CFTR gene. 3. Patient age ≥ 2 years. 4. Provider justification of necessity of medication change if currently stable on another CF regimen and asymptomatic. 5. Patient has not undergone an organ transplant. 6. Medication ordered by Pulmonologist. 7. Approval Duration: 12 months 	<ol style="list-style-type: none"> 1. Provider attestation of continued benefit without adverse drug effects. 2. Provider attestation of continued monitoring as appropriate. 3. Renewal Duration: 12 months.
lumasiran (Oxlumo) injection 94.5mg/0.5ml	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: 	<ol style="list-style-type: none"> 1. All initial approval criteria is met, and 2. Submission of medical records documenting a positive clinical

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> • treatment of primary hyperoxaluria type 1 (PH1) to lower urinary and plasma oxalate levels in pediatric and adult patients. <ol style="list-style-type: none"> 2. Diagnosis of primary hyperoxaluria type 1 (PH1) confirmed by documentation of genetic test results showing a mutation in the alanine:glyoxylate aminotransferase (AGXT) gene OR liver enzyme analysis demonstrating absent or significantly reduced alanine: glyoxylate aminotransferase (AGT) activity. 3. Metabolic testing demonstrating one of the following: <ul style="list-style-type: none"> • Increased urinary oxalate excretion (e.g., > 1 mm/1,73 m² per day [90 mg/1.73 m²]), increased urinary oxalate: creatinine ratio relative to normative values for age OR • Increased plasma oxalate and glyoxylate concentrations. 4. Patient has not received a liver transplant. 5. Prescribed by or in consultation with a nephrologist or other provider (i.e., geneticist, urologist) with experience in treating PH1. 6. Approval Duration: 3 months. 	<p>response to therapy from pre-treatment baseline.</p> <ol style="list-style-type: none"> 3. Approval Duration: 3 months.
lumateperone (Caplyta) capsules 10.5mg, 21mg, 42mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Treatment of depressive episodes associated with bipolar disorder I or II in adults as monotherapy or as an adjunct to lithium or valproate. • Treatment of schizophrenia in adults. 2. Documented trial and failure of at least two other antipsychotic medications indicated to treat the medical diagnosis. <ul style="list-style-type: none"> • Bipolar depression: lurasidone, olanzapine, quetiapine, or risperidone 	<p>Limitations of use:</p> <ul style="list-style-type: none"> • Caplyta is not approved for the treatment of patients with dementia-related psychosis and will not be approved for this indication. • Use with caution in patients at risk of seizures or with conditions that lower the seizure threshold.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> • Schizophrenia: aripiprazole, lurasidone, olanzapine, paliperidone, quetiapine, risperidone, ziprasidone <ol style="list-style-type: none"> 3. Risk versus benefit evaluation if being ordered for adults older than 65 years. 4. Medication ordered by a psychiatrist or other behavioral health specialist. 	
lurasidone (Latuda) tablets 20mg, 40mg, 60mg, 80mg, 120mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Schizophrenia in adults and adolescents (13 to 17 years). • Depressive episode associated with Bipolar I Disorder (bipolar depression) in adults and pediatric patients (10 to 17 years) as monotherapy. • Depressive episode associated with Bipolar I Disorder (bipolar depression) in adults as adjunctive therapy with lithium or valproate. 2. Medication ordered by psychiatrist or other behavioral health specialist. 	Limitations of use: <ul style="list-style-type: none"> • Lurasidone is not approved for the treatment of patients with dementia-related psychosis and will not be approved for this indication.
lusutrombopag (Mulpleta) tablets 3mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Treatment of thrombocytopenia in adult patients with chronic liver disease who are scheduled to undergo a procedure. 2. Patient age \geq 18 years. 3. Not being ordered for patient with chronic liver disease to normalize platelet counts. 4. Dose: 3 mg (1 tablet) daily for 7 days. 5. Approval Duration: one treatment course. 	Each treatment course requires a separate PA request. Initial criteria applies to all requests.
macitentan (Opsumit) 10 mg tablets	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Treatment of pulmonary arterial hypertension (PAH) (WHO Group I) to reduce risks of disease progression and hospitalization. 	<ol style="list-style-type: none"> 1. Documentation of positive clinical response. 2. Approval Duration: 12 months.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ol style="list-style-type: none"> 2. Patient age \geq 18 years. 3. Patient has WHO functional class II-IV PAH, or is class I and at high risk for disease progression. 4. Diagnosis is confirmed by right heart catheterization. 5. Patient has completed a previous at least 4-week trial of ambrisentan OR has a contraindication or intolerance to using it. 6. Patient is not concurrently prescribed another endothelin receptor agonist (ambrisentan, bosentan) 7. NOTE: if the patient is also prescribed for a PDE-5 inhibitor medication (tadalafil, sildenafil), please redirect to Opsynvi. 8. Prescribed by or in consultation with a cardiologist, pulmonologist, or rheumatologist. 9. Quantity Limits: 30 tablets per 30 days 10. Approval Duration: 12 months. 	
macitentan and tadalafil (Opsynvi) 10-20 mg, 10-40 mg tablets	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Treatment of chronic pulmonary arterial hypertension (PAH, WHO Group I) in adult patients of WHO functional class II-III. 2. Patient age \geq 18 years. 3. Patient has WHO functional class II-IV PAH, or is class I and at high risk for disease progression. 4. Patient is not concurrently prescribed another endothelin receptor agonist (macitentan, ambrisentan, bosentan), PDE-5 inhibitor (sildenafil, tadalafil), or Adempas (riociguat). 5. Diagnosis is confirmed by right heart catheterization. 6. Prescribed by or in consultation with a cardiologist, pulmonologist, or rheumatologist. 7. Quantity Limits: 30 tablets per 30 days 8. Approval Duration: 12 months. 	<ol style="list-style-type: none"> 1. Documentation of positive clinical response to therapy. 2. Approval Duration: 12 months.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
maribavir (Livtency) tablets 200mg STEP THERAPY	<ol style="list-style-type: none"> Ordered for an approved indication for use: <ul style="list-style-type: none"> treatment of adults and pediatric patients (12 years of age and older and weighing at least 35 kg) with post-transplant CMV infection/disease that is refractory to treatment (with or without genotypic resistance) with ganciclovir, valganciclovir, cidofovir or foscarnet. Medication is not prescribed in conjunction with ganciclovir or valganciclovir. Medication is prescribed by or in consultation with a hematologist, infectious disease specialist, oncologist or physician affiliated with a transplant center. Approval Duration: not to exceed 8 weeks. 	If a patient has a paid claim in the MFC system for ganciclovir, valganciclovir, cidofovir, or foscarnet, Livtency will process at the pharmacy without PA. If there is no evidence of a paid claim for ganciclovir, valganciclovir, cidofovir, or foscarnet, a PA is required, and documentation of previous use of one of these medications should be submitted.
mepolizumab (Nucala) injection 40mg/0.4mL syringes 100mg pens, syringes	<ol style="list-style-type: none"> Ordered for an approved indication for use: <ul style="list-style-type: none"> Add-on maintenance treatment for severe asthma with eosinophilic phenotype in patients aged 6 years and older. Add-on treatment of adult patients with chronic rhinosinusitis with nasal polyps. Treatment of eosinophilic granulomatosis with polyangiitis (EGPA) in adults. Treatment of adult and pediatric patients aged \geq 12 years of age with hypereosinophilic syndrome (HES) for \geq 6 months without an identifiable non-hematologic secondary cause. Approval is indication specific: Asthma: <ul style="list-style-type: none"> Patient is \geq 6 years or age; AND Patient has blood eosinophil level \geq 150 cells/μL within previous 6 weeks or within 6 weeks prior to treatment with Nucala or another monoclonal 	Asthma: <ul style="list-style-type: none"> Patient has already received 6 months of therapy with Nucala. Patient continues to receive therapy with one inhaled corticosteroid or one inhaled corticosteroid-containing combination inhaler. Patient has responded to therapy (e.g. decreased asthma exacerbations, symptoms, hospitalizations, ER visits, urgent care visits, or decreased requirement for oral corticosteroid therapy. Chronic Rhinosinusitis with Nasal Polyps:

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>antibody therapy that reduces blood eosinophil levels, (e.g. Cinqair, Dupixent, Fasenna, Nucala, Tezspire or Xolair); AND</p> <ul style="list-style-type: none"> • Patient has received at least three consecutive months of combination therapy with BOTH an inhaled corticosteroid AND at least one additional asthma controller or asthma maintenance medication; AND • Patient has asthma that is controlled or was uncontrolled at baseline as defined by one of the following: <ul style="list-style-type: none"> ○ Patient experienced 2 or more asthma exacerbations requiring treatment with systemic corticosteroids in the previous year; OR ○ Patient experienced at least one asthma exacerbation requiring hospitalization, an emergency department visit, or urgent care visit in the previous year; OR ○ Patient has a forced expiratory volume in 1 second (FEV₁) < 80% predicted; or ○ Patient has an FEV₁/forced vital capacity (FVC) < 0.80; OR ○ Patient has asthma that worsens upon tapering of oral (systemic) corticosteroid therapy. • Medication ordered by an Allergist, Immunologist or Pulmonologist. • Approval Duration: 6 months. <p><u>Chronic Rhinosinusitis with Nasal Polyps:</u></p> <ul style="list-style-type: none"> • Patient is ≥ 18 years of age; and 	<ul style="list-style-type: none"> • Patient has received at least 6 months of therapy with Nucala. • Patient continues to receive therapy with an intranasal corticosteroid; and • Patient has responded to therapy (e.g. reduced nasal polyp size, improved nasal congestion, reduced sinus opacification, decreased sino-nasal symptoms, improved sense of smell.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> • Patient has chronic rhinosinusitis with nasal polyps as evidenced by direct examination, endoscopy, or sinus computed tomography (CT) scan; and • Has had two or more of the following symptoms for at least 6 months: nasal congestion, nasal obstruction, nasal discharge, and/or reduction or loss of smell; AND • Patient has received at least 4 weeks of therapy with an intranasal corticosteroid; AND • Patient will continue to receive therapy with an intranasal corticosteroid concomitantly with Nucala; and • Patient meets one of the following: <ul style="list-style-type: none"> ○ Patient has had at least one course of treatment with systemic corticosteroid for 5 days or more within the previous 2 years; or ○ Patient has a contraindication to systemic corticosteroid therapy, or ○ Patient has prior history of surgery for nasal polyps; AND • Prescribed by or in consultation with an allergist, immunologist, or otolaryngologist/ENT. • Approval Duration: 6 months. <p>3. Indications excluded from coverage include:</p> <ul style="list-style-type: none"> • Atopic Dermatitis • COPD • Concurrent use of another monoclonal antibody therapy. • Eosinophilic esophagitis, eosinophilic gastroenteritis, or eosinophilic colitis 	

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
methadone (for pain) solution 5mg/5ml, 10mg/5ml tablets 5mg, 10mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • The management of chronic pain severe enough to require daily, around-the- clock, long-term opioid treatment and for which alternative treatment options are inadequate. 2. Completion of an opioid prior authorization form. 3. Submission of clinical documentation from last office visit, dated within 3 months of the request. 4. Maximum approval duration is 6 months but may be reduced based on any of the criteria as outlined in Pharmacy Policy 219.DC: Opioid Prescription Prior Authorization. 	<p>All long-acting opioids require Prior Authorization (PA). The PA form can be accessed using the following link:</p> <p>OPIOID PRIOR AUTH FORM-DC</p>
methylphenidate (Jornay PM) extended-release capsules	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • To treat symptoms of attention deficit hyperactivity disorder (ADHD) in children ages 6 years and older and adults 2. Step therapy: at least 4-week trial of an amphetamine salt combination AND a 4-week trial of methylphenidate. 3. Patient has documented need for evening medication administration (e.g. frequent missed morning doses, autism diagnosis) 4. Will not be used concurrently with other stimulant medications. Jornay is designed with an extendable duration of effect and can be dose adjusted to increase the duration of symptom control time. 5. Prescribed by or in conjunction with a psychiatrist or other behavioral health specialist. 6. Approval Duration: 12 months 	<ol style="list-style-type: none"> 1. Initial criteria continue to be met. 2. Patient has documented positive clinical response with the medication since starting the medication. 3. Approval duration: 12 months
mifepristone (Korlym) tablets Korlym-300mg ONLY	<ol style="list-style-type: none"> 7. Ordered for an approved indication for use: 	<ol style="list-style-type: none"> 4. Documentation of one of the following:

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> • Control of hyperglycemia secondary to hypercortisolism in adult patients with endogenous Cushing’s syndrome who have type 2 diabetes mellitus or glucose intolerance and have failed surgery or are not candidates for surgery. <ol style="list-style-type: none"> 8. Patient is ≥ 18 years of age. 9. Diagnosis of endogenous Cushing’s syndrome (i.e., hypercortisolism is not a result of chronic administration of high dose glucocorticoids). 10. Patient has type 2 diabetes mellitus OR the patient has glucose intolerance as defined by a 2-hour glucose tolerance test value of 140-199 mg/dL. 11. The patient has either failed surgery or is not a candidate for pituitary surgery. 12. Prescribed by or in consultation with an endocrinologist. 13. The dose does not exceed 20 mg/kg/day. 14. Approval Duration: 3 months. 	<ul style="list-style-type: none"> • Patient has improved glucose tolerance while on Korlym therapy; or • Patient has stable glucose tolerance while on Korlym therapy. <ol style="list-style-type: none"> 5. Dose does not exceed 20 mg/kg/day. 6. Approval duration: 3 months per authorization. <p>**No other indications approved, can redirect requests for Mifeprex brand to appropriate formulary alternatives.</p>
mirikizumab (Omvo) injection 100 mg/1 ml	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Maintenance treatment of ulcerative colitis (UC) in adults with moderate to severe active disease. 1. Patient is ≥ 18 years of age, and 2. Patient has had a trial of one systemic agent for UC (e.g., 6-MP, azathioprine, cyclosporine, tacrolimus or a corticosteroid. Note that trial of a mesalamine product does <u>not</u> count as a systemic therapy for UC) OR 3. Patient has both: <ul style="list-style-type: none"> • Pouchitis AND • Patient has tried an antibiotic, probiotic, corticosteroid enema, or mesalamine enema 4. Patient is not being treated concurrently with a 	<ol style="list-style-type: none"> 1. Patient exhibits a positive clinical response by at least one objective measure from baseline. (e.g., fecal calprotectin levels, C-reactive protein, endoscopic assessment, and/or decreased utilization of corticosteroids OR 2. Patient has a documented clinical improvement in at least one subjective measure from baseline (e.g., decreased pain, fatigue, stool frequency, and/or rectal bleeding). 3. Approval duration: 12 months.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>biologic or targeted synthetic disease-modifying antirheumatic drug (DMARD) for UC. (e.g., adalimumab, infliximab, sarilumab, abatacept, rituximab, ustekinumab, apremilast, ozanimod, or similar).</p> <p>5. Medication is prescribed by or in consultation with a gastroenterologist.</p> <p>6. Initial Approval Duration: 6 months; if patient has already received > 6 months of subcutaneous therapy, then approval duration is 12 months.</p>	
mitapivat (Pyrukynd) tablets 5mg, 20mg, 50mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • The treatment of hemolytic anemia in adults with pyruvate kinase (PK) deficiency 2. Confirmatory genetic testing of PKLR gene showing \geq 2 variant alleles with at least one- missense mutation in the liver and red blood cell (PKLR) gene. 3. Patient is not homozygous for the c.1436G>A (p.R479H) variant. 4. Patient does not have two non-missense variants (without the presence of another missense variant) in the PKLR gene. 5. Baseline hemoglobin less than or equal to 10 g/dL. 6. Prescribed by or in consultation with a Hematologist. 7. Initial Approval Duration limited to 6 months. 	<ol style="list-style-type: none"> 1. Documentation of positive clinical response to Pyrukynd therapy based on ONE of the following: <ul style="list-style-type: none"> • Patient has been on Pyrukynd for > 52 weeks and has maintained positive clinical response to therapy; OR • Reduction in transfusions of \geq 33% in the number of red blood cell units transfused during the initial 24-week period compared with the patient’s historical transfusion burden; OR • A \geq 1.5 g/dL increase in hemoglobin from baseline sustained at 2 or more scheduled assessments 4 weeks apart during the initial 24-week period without any transfusions. 2. Authorization duration: 12 months <ul style="list-style-type: none"> • If documentation does not provide evidence of positive clinical response to Pyrukynd therapy, allow for dose titration with discontinuation of therapy. In this

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
modafinil (Provigil) tablets 100mg, 200mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • to improve wakefulness in adult patients with excessive sleepiness associated with narcolepsy, obstructive sleep apnea, or shift work disorder. 2. Will not be used concurrently with monoamine oxidase inhibitors (isocarboxazid, phenylzine, tranylcypromine) 3. No past medical history of left ventricular hypertrophy 4. If PMH of mitral valve prolapse, no previous occurrence of mitral valve prolapse syndrome with past use of CNS stimulants (including but not limited to ischemic ECG changes, chest pain, or arrhythmia) 5. No active symptoms of mania or psychosis 6. If the patient has a history of cardiovascular disease, psychosis, depression or mania, patient is on a treatment regimen with adequate disease state control, AND prescriber has conducted a risk vs. benefit assessment to ensure clinical appropriateness before modafinil initiation 7. Confirmation of negative pregnancy status within 1 week of treatment initiation for females of reproductive age. 8. Approval Duration: 12 months 	<p>case, authorization duration is for 4 weeks.</p> <ul style="list-style-type: none"> • Maximum recommended daily dose is 200 mg (single dose or 2x100 mg doses); evidence is limited for daily doses >200 mg regardless of indication <p><u>Renewal criteria:</u></p> <ul style="list-style-type: none"> • Documentation of positive clinical response with modafinil treatment • Initial approval criteria continue to be met • Renewal duration: 12 months
morphine sulfate extended-release (MS Contin) tablets 15mg, 30mg, 60mg 100mg, 200mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • The management of pain severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate. 2. Completion of an opioid prior authorization form. 	<p>All long-acting opioids require Prior Authorization (PA). The PA form can be accessed using the following link:</p> <p>OPIOID PRIOR AUTH FORM-DC</p>

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ol style="list-style-type: none"> 3. Submission of clinical documentation from last office visit, dated within 3 months of the request. 4. Maximum approval duration is 6 months but may be reduced or denied based on the criteria as outlined in Pharmacy Policy 219.DC: Opioid Prescription Prior Authorization. 	
nintedanib (Ofev) capsule 100mg, 150mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Treatment of adults for idiopathic pulmonary fibrosis. • Treatment of adults for chronic fibrosing interstitial lung diseases (ILDs) with a progressive phenotype. • To slow the rate of decline in pulmonary function in patients with systemic sclerosis associated interstitial lung disease (SSc-ILD). 2. Documentation that patient does not smoke. 3. Medication ordered by a pulmonologist. 4. Authorization Duration: 12 months. 	<ol style="list-style-type: none"> 1. All initial criteria are met. 2. Documentation of positive clinical response to Ofev therapy. 3. Approval Duration: 12 months
nirogacestat (Ogsiveo) tablets 150 mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Treatment of Desmoid Tumors (aggressive fibromatosis) in adult patients. 2. Patient is aged 18 years or older. 3. Patient has been diagnosed with progressing desmoid tumors as defined as $\geq 20\%$ progression within 12 months, AND 4. The desmoid tumors are not amenable to surgery or radiotherapy, AND 5. The patient requires systemic treatment. 6. Quantity Limits: 2 tablets daily, not to exceed 150 mg BID. 7. Approval Duration: 12 months. 	<ol style="list-style-type: none"> 1. Patient does not show evidence of progressive disease while on Ogsiveo therapy. 2. Approval Duration: 12 months
nitisinone (Orfadin) capsules	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: 	<ol style="list-style-type: none"> 1. Meets all initial approval criteria.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
2mg, 5mg, 10mg, 20mg	<ul style="list-style-type: none"> • treatment of adult and pediatric patients with hereditary tyrosinemia type 1 (HT-1) in combination with dietary restriction of tyrosine and phenylalanine. <ol style="list-style-type: none"> 2. Diagnosis of type 1 tyrosinemia by biochemical or DNA testing. 3. Patient adherent to dietary restrictions of tyrosine and phenylalanine. 4. Patient is under the care of a nutritionist. 5. Dose not to exceed 2 mg/kg/day. 6. Patient is not enrolled in any study involving the requested drug. 7. PA form completed completely. 8. Approval Duration: 3 months 	<ol style="list-style-type: none"> 2. MDH provided PA form completed with all required documentation. 3. Approval duration: 3 months.
nusinersen (Spinraza)	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Diagnosis of SMA Type I, II, or III. • Diagnosis by a neurologist with expertise in the diagnosis of SMA; 2. Genetic testing confirming both: <ul style="list-style-type: none"> • 5q SMA homozygous gene deletion, homozygous gene mutation, or compound heterozygous mutation: AND • At least 2 copies of SMN2 3. AND <ul style="list-style-type: none"> • Patient is not dependent on invasive ventilation or tracheostomy. • Patient is not dependent on non-invasive ventilation beyond use for naps and nighttime sleep; • Patients with Type II and III SMA must have some functional upper extremity use. 4. Initial therapy 	<ol style="list-style-type: none"> 1. Cannot be used in combination with Zolgensma (onasemnogene abeparvovec). 2. Each Spinraza maintenance dose must be preauthorized; 3. Approval period is 3 months. 4. All the criteria for initial therapy must be met: <ul style="list-style-type: none"> • Medical records must be submitted that document repeat motor testing since the most recent Spinraza® dose using the same motor test done to establish baseline motor ability, unless it is determined that the original test is no longer appropriate;

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> • Medical records must be submitted documenting all of the above criteria; • Medical records must be submitted documenting a baseline motor examination utilizing at least one of the following exams (based on patient age and motor ability) to establish baseline motor ability. • Hammersmith infant neurological exam (HINE); • Hammersmith Functional Motor Scale Expanded (HFMSE); • Upper Limb Module Test (non-ambulatory; or • Childrens Hospital of Philadelphia Infant Test of Neuromuscular Disorders (CHOP-INTEND) <ol style="list-style-type: none"> 5. Prescribed by a neurologist with expertise in the treatment of SMA. 6. Spinraza must be given according to the current FDA labelling guidelines for dosage and timing; 7. Must be administered intrathecally by a physician or other healthcare professional experienced in performing lumbar punctures. 8. Initial and Renewal Approval Duration: 4 loading doses for initiation, 3 months each approval period thereafter. 	<ul style="list-style-type: none"> • Repeat motor testing must document a response to treatment as defined by the following: <p>HINE:</p> <ol style="list-style-type: none"> 1. Improvement or maintenance of previous improvement of at least 2 points (or max score of 4) in ability to kick (improvement in at least 2 milestones); OR 2. Improvement or maintenance of previous improvement of at least 1 point increase in motor milestones of head control, rolling, sitting, crawling, standing, or walking (consistent with improvement by at least 1 milestone); AND 3. Improvement or maintenance of previous improvement in more HINE motor milestones. <p>HFMSE:</p> <ol style="list-style-type: none"> 1. Improvement or maintenance of improvement of at least a 3-point increase in score; <p>ULM:</p> <ol style="list-style-type: none"> 1. Improvement or maintenance of previous improvement of at least 2-point increase in score; <p>CHOP-INTEND:</p> <ol style="list-style-type: none"> 1. Improvement or maintenance of previous improvement of at least a 4-point increase in score.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
<p>ocrelizumab (Ocrevus) IV injection 300mg/10ml</p> <p>ocrelizumab and hyaluronidase (Ocrevus Zunovo) SQ injection 920-23,000 mg-units/23mL</p>	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Primary progressive multiple sclerosis (MS); • Relapsing forms of MS, to include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease. 2. Age is ≥18 years and <55 years of age. 3. Patient has one of the following: <ul style="list-style-type: none"> • Ineffective treatment response due to continued clinical relapse, intolerance, or contraindication to two or more MS drugs; • Patient is not a candidate for any other preferred first-line treatments due to MS severity; • Patient is at higher risk of poor long-term outcome (spinal cord involvement, highly active disease, poor relapse recovery), as determined by their neurologist. 4. Not being used in combination with other immunomodulating or immunosuppressive therapies, including immunosuppressant doses of corticosteroids. 5. Not being used in combination with another MS disease modifying agent [Avonex, Betaseron, dimethyl fumarate, Extavia, fingolimod, glatiramer, glatopa, Kesimpta, Mayzent, Rebif, teriflunomide, Vumerity]. 6. Medication ordered by a neurologist. 7. Approval duration: 12 months. 	<ol style="list-style-type: none"> 1. All initial criteria continue to be met. 2. Documentation of positive clinical response to Ocrevus therapy. 3. Approval duration: 12 months.
<p>olanzapine and samidorphan (Lybalvi) tablets 5mg/10mg, 10mg/10mg, 15mg/10mg, 20mg/10mg</p>	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Schizophrenia in adults • Bipolar I disorder in adults as acute treatment of manic or mixed episodes as monotherapy and as adjunct to lithium or valproate 	<ol style="list-style-type: none"> 1. All initial criteria continue to be met. 2. Documentation of positive clinical response to therapy. 3. Approval duration: 12 months.

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> • Bipolar I disorder in adults as maintenance monotherapy treatment <ol style="list-style-type: none"> 2. Clinical documentation the patient does not have active opioid use disorder, AND 3. Patient is not concurrently prescribed opioid medication(s), AND 4. Patient is not actively going through opioid withdrawal, AND 5. Negative baseline urine drug screen result from within the last 14 days. 6. Patient has a history of failure, contraindication, or intolerance to at least THREE (3) formulary atypical antipsychotic agents, such as: <ul style="list-style-type: none"> • aripiprazole • olanzapine • quetiapine IR or XR • risperidone • ziprasidone 7. Approval Duration: 12 months 	
olipudase alfa-rpcp (Xenpozyme) 4mg, 20mg	<ol style="list-style-type: none"> 1. Prescribed for an approved indication for use; treatment of non-central nervous system manifestations of acid sphingomyelinase deficiency (ASMD) in adult and pediatric patients. 2. Acid sphingomyelinase enzyme assay (as measured in peripheral leukocytes, cultured fibroblasts, or lymphocytes) or genetic testing results documenting a mutation in the sphingomyelin phosphodiesterase-1 (SMPD1) gene. 3. Prior to initiation of Xenpozyme, baseline transaminase (alanine aminotransferase [ALT] and aspartate aminotransferase [AST]) levels recorded within past 30 days. 	<ol style="list-style-type: none"> 1. Documented response to therapy improvement or stabilization in disease (e.g., improvement in lung function, reduction in spleen volume, reduction in liver volume, improvement in platelet count, improvement in linear growth progression). 2. Documentation of patient's current weight. 3. Dose does not exceed 3 mg/kg IV every 2 weeks. 4. Approval duration: 3 months

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ol style="list-style-type: none"> 4. Documentation of patient's current weight. 5. Prescribed by or in consultation with a metabolic disease specialist or geneticist. 6. Dose does not exceed 3 mg/kg IV every 2 weeks. 7. Approval duration: 3 months 	
<p>omalizumab (Xolair) Injection 75mg/0.5ml, 150mg/ml Solution for injection 150mg</p>	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • moderate to severe persistent asthma in patients \geq 6 years of age with a positive skin test or in vitro reactivity to a perennial aeroallergen and symptoms that are inadequately controlled with inhaled corticosteroids. • chronic spontaneous urticaria (CSU) in adults and adolescents \geq 12 years of age who remain symptomatic despite H1 antihistamine treatment. • Chronic rhinosinusitis with nasal polyps (CRSwNP) in adult patients \geq 18 years of age with inadequate response to nasal corticosteroids, as add-on maintenance treatment. • IgE-mediated food allergy and patients \geq 1 year of age for the reduction of Type I allergic reactions, including anaphylaxis, that may occur with accidental exposure to one or more foods, in conjunction with food allergen avoidance. 2. NOT eligible for coverage for the treatment/management of: <ul style="list-style-type: none"> • Acute bronchospasm or status asthmaticus, or • Emergency treatment of allergic reactions, including anaphylaxis, or • Other forms of urticaria. 3. Patient is not receiving treatment in combination with ANY of the following: <ul style="list-style-type: none"> • Anti-interleukin-4 therapy (e.g. Dupixent 	<p>Renewal criteria applicable to all indications in addition to indication specific criteria outlined below:</p> <ol style="list-style-type: none"> 1. Patient is not receiving treatment in combination with ANY of the following: <ul style="list-style-type: none"> • Anti-interleukin-5 therapy (e.g., Cinqair (reslizumab), Fasenna (benralizumab), Nucala (mepolizumab)). • Anti-interleukin-4 therapy (e.g., Dupixent (dupilumab)). • Thymic stromal lymphopoietin (TSLP) inhibitor (e.g., Tezspire, (Tezepelumab)). <p>Asthma:</p> <ol style="list-style-type: none"> 1. Documentation of positive clinical response (e.g. reduction in frequency of exacerbations, decreased use of rescue medications, increase in percent predicted FEV1 from pre-treatment baseline or reduction in symptom severity or frequency), AND 2. Xolair is being used in combination with an ICS-containing maintenance

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>(dupilumab)).</p> <ul style="list-style-type: none"> • Anti-interleukin-5 therapy (e.g., Cinqair (reslizumab), Fasentra, (benralizumab), Nucala (mepolizumab)). • Thymic stromal lymphopoietin (TSLP) inhibitor (e.g., Tezspire (Tezepelumab)). <p>Asthma:</p> <ol style="list-style-type: none"> 1. Patient aged ≥ 6 years of age. 2. Positive skin test or in-vitro reactivity to a perennial aeroallergen, AND 3. Submission of clinical documentation showing the baseline (pre-treatment) serum total IgE level ≥ 30 IU/ml and ≤ 1300 IU/ml, AND 4. Diagnosed with moderate to severe asthma inadequately controlled with inhaled corticosteroids as defined by at least ONE of the following: <ul style="list-style-type: none"> • Poor symptom control (e.g., Asthma Control Questionnaire (ACQ) score consistently greater than 1.5 or Asthma Control Test (ACT) score consistently less than 20); or • Two or more bursts of systemic corticosteroids for at least 3 days each in previous 12 months; or • Asthma-related emergency treatment (ER visit, hospital admission, or unscheduled OV for nebulizer or emergency treatment); OR • Airflow limitation (e.g., after appropriate bronchodilator withhold forced expiratory volume in 1 second (FEV1) less than 80% predicted; OR • Patient is currently dependent on oral corticosteroids for the treatment of asthma; AND, 5. Xolair will be used in combination with one maximally dosed combination ICS/LABA inhaler OR with an ICE 	<p>medication – NOT covered as monotherapy.</p> <ol style="list-style-type: none"> 3. Approval Duration: 12 months. <p><u>IgE-mediated Food Allergy:</u></p> <ol style="list-style-type: none"> 1. Documentation of positive clinical response to Xolair therapy, e.g. reduction in type I allergic reactions, and 2. Used in conjunction with food allergen avoidance, and 3. Patient has access to epinephrine, and 4. Prescribed by an allergist or immunologist. 5. Approval Duration: 12 months. <p><u>Rhinosinusitis, chronic, with nasal polyps (CRSwNP):</u></p> <ol style="list-style-type: none"> 1. Documentation of positive clinical response to Xolair therapy. 2. Patient continues to use add-on maintenance therapy with intranasal corticosteroids – NOT covered as monotherapy. 3. Approval Duration: 12 months. <p><u>Urticaria (chronic spontaneous):</u></p> <ol style="list-style-type: none"> 1. Documentation of positive clinical response to Xolair therapy (e.g. reduction in exacerbations, itch severity, hives). 2. Approval Duration: 12 months.

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>inhaler and one additional asthma controller medication (e.g. montelukast, theophylline).</p> <p>6. Prescribed by an allergist, immunologist or pulmonologist.</p> <p>7. Approval Duration: 12 months.</p> <p><u>IgE-mediated Food Allergy:</u></p> <ol style="list-style-type: none"> 1. Patient aged ≥ 1 year of age. 2. Diagnosis of IgE-mediated food allergy to one or more foods, AND 3. Diagnosis has been confirmed by BOTH of the following: <ul style="list-style-type: none"> • History of type I allergic reactions (e.g., nausea, vomiting, cramping, diarrhea, flushing, pruritus, urticaria, swelling of lips, face, or throat, wheezing, lightheadedness, syncope), AND • ONE of the following: <ul style="list-style-type: none"> ○ Food specific skin prick testing (SPT) ○ IgE antibody in vitro testing ○ Oral food challenge (OFC) 4. Xolair will be used in conjunction with food allergen avoidance, AND 5. Patient has access to epinephrine, AND 6. Prescribed by an allergist or immunologist. 7. Approval Duration: 12 months. <p><u>Rhinosinusitis, chronic, with nasal polyps (CRSwNP):</u></p> <ol style="list-style-type: none"> 1. Patient aged ≥ 18 years of age. 2. Prescribed as add-on maintenance treatment to nasal corticosteroids (NOT covered as monotherapy). 3. Diagnosis of nasal polyps, AND 4. Patient has TWO or more of the following symptoms for ≥ 12 weeks: 	

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> • Nasal mucopurulent discharge • Nasal obstruction, blockage, or congestion • Facial pain, pressure and/or fullness • Reduction or loss of sense of smell; AND <p>5. ONE of the following findings using nasal endoscopy and/or sinus computed tomography (CT):</p> <ul style="list-style-type: none"> • Purulent mucus or edema in the middle meatus or ethmoid regions, or • Polyps in the nasal cavity or the middle meatus, or • Radiographic imaging demonstrating mucosal thickening or partial or complete opacification of paranasal sinuses; AND <p>6. ONE of the following:</p> <ul style="list-style-type: none"> • Patient has not obtained relief after a trial of BOTH intranasal corticosteroids and one other therapy used in the management of nasal polyps (e.g. nasal saline irrigations, antileukotriene agents); OR • Patient has required systemic corticosteroids for nasal polyps in the previous 2 years; OR • Patient has required prior sinus surgery, AND <p>7. Patient will receive Xolair as add-on maintenance therapy in combination with intranasal corticosteroids.</p> <p>8. Prescribed by an allergist, immunologist, otolaryngologist, or pulmonologist.</p> <p>9. Approval Duration: 12 months.</p> <p><u>Urticaria (chronic spontaneous):</u></p> <ol style="list-style-type: none"> 1. Patient is aged \geq 12 years of age. 2. Patient remains symptomatic following: 	

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> • at least a 2-week trial of, contraindication, or intolerance to TWO H1-antihistamines (e.g. fexofenadine, diphenhydramine, loratadine OR • a two-week trial of taking a second-generation H1-antihistamines in combination with: <ul style="list-style-type: none"> ○ a different second generation H1 antihistamine, or ○ a first generation H1 antihistamine (e.g. hydroxyzine, diphenhydramine, or chlorpheniramine), or ○ an H2 antihistamine (e.g. famotidine or cimetidine), or ○ a leukotriene modifier (e.g. montelukast). <p>3. Prescribed by an allergist, dermatologist or immunologist</p> <p>4. Approval Duration: 12 months.</p>	
omega-3-acid ethyl esters (Lovaza) capsules 1 Gram	<p>1. Ordered for an approved indication for use:</p> <ul style="list-style-type: none"> • as an adjunct to diet to reduce triglyceride levels in adult patients with severe (≥ 500 mg/dL) hypertriglyceridemia <p>2. Patient is ≥ 18 years old.</p> <p>3. Member must have tried and failed a 30-day trial of OTC fish oil.</p> <p>4. Approval Duration: 12 months.</p>	<p>1. The patient has achieved or maintained a reduction in triglyceride levels from baseline.</p> <p>2. Approval Duration: 12 months.</p>
onabotulinumtoxinA (Botox) injection 100 Unit, 200 Unit	<p>1. Ordered for an approved indication for use:</p> <ul style="list-style-type: none"> • Overactive bladder (OAB) with symptoms of urge urinary incontinence, urgency, and frequency, in adults who have an inadequate response to or are intolerant of an anticholinergic medication. • Urinary incontinence due to detrusor overactivity associated with a neurologic condition [e.g., 	<p>Limitations for Use:</p> <ul style="list-style-type: none"> • Botox will NOT be approved for cosmetic purposes • Safety and effectiveness have <u>not</u> been established for:

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>spinal cord injury, multiple sclerosis] in adults who have an inadequate response to or are intolerant of an anticholinergic medication.</p> <ul style="list-style-type: none"> • Neurogenic detrusor overactivity (NDO) in pediatric patients ≥ 5 years of age who have an inadequate response to or are intolerant of anticholinergic medication. • Prophylaxis of headaches in adult patients with chronic migraine (≥15 days per month with headache lasting ≥ 4 hours a day). • Spasticity in adult patients. • Cervical dystonia in adult patients to reduce the severity of abnormal head position and neck pain. • Severe axillary hyperhidrosis of adults inadequately managed by topical agents. • Treatment of blepharospasm associated with dystonia in patients 12 years of age and older. • Treatment of strabismus in patients 12 years of age and older. <p>2. Medication ordered by a Neurologist, Urologist, Ophthalmologist, or applicable specialist.</p>	<ul style="list-style-type: none"> • Prophylaxis of episodic migraine (≤ 14 headache days/month). • treatment of upper or lower limb spasticity in pediatric patients.
<p>Opioids</p>	<p>FOR IMPORTANT INFORMATION ABOUT PRESCRIBING OPIOIDS FOR MEDSTAR FAMILY CHOICE MEMBERS, PLEASE VISIT THE OPIOID PRIOR AUTHORIZATION REQUIREMENTS PAGE OF THE MFC-DC WEBSITE.</p> <ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • The management of pain severe enough to require opioid treatment and for which alternative treatment options are inadequate. 2. Completion of the opioid prior authorization form. 	<p>The Opioid PA form can be accessed using the following link:</p> <p>OPIOID PRIOR AUTH FORM-DC</p>

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ol style="list-style-type: none"> 3. Submission of supporting clinical documentation for the last office visit, dated within the previous 3 months. 4. Maximum approval duration is 6 months but may be approved for a shorter duration based on any of the criteria as outlined in Pharmacy Policy 219.DC: Opioid Prescription Prior Authorization. 	
oxcarbazepine extended release 24-hour (Oxtellar XR) tablets 150mg, 300mg, 600mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Treatment of partial-onset seizures in adults and in children ≥ 6 years of age. 2. Treatment failure, adverse effects, or contraindication to formulary preferred agents. 3. Medication ordered by a Neurologist. 	Immediate-release and extended-release preparations are not bioequivalent and not interchangeable on a mg per mg basis.
oxycodone IR capsules, tablets, oral solution/concentrate 5 mg capsules 100mg/5mL oral concentrate 5mg/5mL oral solution IR tablets 5, 10, 13, 20, 30 mg Oxycontin ER tablets 10, 15, 20, 30, 40 mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • The management of pain severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate. 2. Completion of the opioid prior authorization form. 3. Submission of supporting clinical documentation for last office visit dated within previous 3 months. 4. Maximum approval duration is 6 months but may be reduced or denied based on any of the criteria as outlined in Pharmacy Policy 219.DC: Opioid Prescription Prior Authorization. 	All long-acting opioids require Prior Authorization (PA). The PA form can be accessed using the following link: OPIOID PRIOR AUTH FORM-DC
oxycodone/acetaminophen tablets, oral solution tablets 5-325, 7.5-325, 10-325 mg oral solution 5-325mg/5mL	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • The management of pain severe enough to require opioid treatment and for which alternative treatment options are inadequate. 2. Completion of the opioid prior authorization form. 3. Submission of supporting clinical documentation for last office visit dated within the previous 3 months. 	All long-acting opioids require Prior Authorization (PA). The PA form can be accessed using the following link: OPIOID PRIOR AUTH FORM-DC

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	4. Maximum approval duration is 6 months but may be reduced or denied based on any of the criteria as outlined in Pharmacy Policy 219.DC: Opioid Prescription Prior Authorization.	
oxycodone extended release 12-hour (Opana) tablets 5mg, 7.5mg, 10mg, 15mg, 20mg, 30mg, 40mg	1. Ordered for an approved indication for use: <ul style="list-style-type: none"> management of pain severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate. 2. Completion of the opioid prior authorization form. 3. Submission of supporting clinical documentation for last office visit, dated within previous 3 months. 4. Maximum approval duration is 6 months but may be reduced or denied based on any of the criteria as outlined in Pharmacy Policy 219.DC: Opioid Prescription Prior Authorization.	All long-acting opioids require Prior Authorization (PA). The PA form can be accessed using the following link: OPIOID PRIOR AUTH FORM-DC
ozanimod (Zeposia) capsules 0.23mg, 0.46mg, and 0.92mg capsules 7-day starter pack Capsule Starter Kit	1. Ordered for an approved indication for use: <ul style="list-style-type: none"> Treatment of relapsing forms of multiple sclerosis (MS), to include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease, in adults. Treatment of moderately to severely active ulcerative colitis (UC) in adults. 2. Patient has not received a manufacturer supplied sample or any form of assistance from the manufacturer coupon or sample card as a means to establish as a current user of Zeposia. 3. Baseline evaluation of the following labs before starting treatment: CBC, ECG, LFT's	Renewal Criteria: 1. Initial approval criteria continue to be met. 2. Patient is not receiving in combination a biologic DMARD or janus kinase inhibitor Multiple Sclerosis: <ul style="list-style-type: none"> Patient experiencing disease stability or improvement while receiving Zeposia. Maximum approval Duration: 12 months Ulcerative Colitis:

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>4. No history (within previous 6 months) of myocardial infarction, unstable angina, stroke, transient ischemic attack, decompensated heart failure.</p> <p>5. No severe untreated sleep apnea</p> <p>6. Zeposia will not be used in combination with either a biologic DMARD (e.g. adalimumab, Simponi (golimumab), Stelara (ustekinumab) OR a Janus kinase inhibitor (e.g. Xeljanz (tofacitinib), Rinvoq (upadacitinib) (Note: Ampyra and Nuedexta are not disease modifying).</p> <p>7. <u>Additional Criteria for Multiple Sclerosis</u></p> <ul style="list-style-type: none"> • Prescribed by or within consultation with a neurologist. <p>8. <u>Additional Criteria for Ulcerative Colitis</u></p> <ul style="list-style-type: none"> • Diagnosis of moderately to severely active UC • Patient has failed, contraindicated or intolerance to a course of oral corticosteroids and/or immunosuppressants (e.g. azathioprine, or 6-mercaptopurine) OR • Patient has been previously treated with a biologic or targeted synthetic DMARD FDA-approved for the treatment of UC as documented by claims history or submission of medical records. (e.g., adalimumab, Simponi (golimumab), Stelara (ustekinumab), Xeljanz (tofacitinib), Rinvoq (upadacitinib)). • Prescribed by or in consultation with a gastroenterologist. <p>9. Approval duration: 12 months</p>	<ul style="list-style-type: none"> • Patient has achieved or maintained remission. • Patient shows positive clinical response as evidenced by low disease activity or improvement in signs/symptoms of the condition when there is improvement in any ONE of the following from baseline: <ul style="list-style-type: none"> ○ Stool frequency ○ Rectal bleeding ○ Urgency of defecation ○ C-reactive protein (CRP) ○ Fecal calprotectin (FC) ○ Endoscopic appearance of the mucosa ○ Improvement on a disease activity scoring tool (e.g. Ulcerative Colitis Endoscopic Index of Severity (UCEIS, Mayo score) <p>3. Approval Duration: 12 months</p>
<p>palbociclib (Ibrance) capsules 75mg, 100mg, 125mg</p>	<p>1. Ordered for an approved indication for use:</p> <ul style="list-style-type: none"> • Treatment of adult patients with hormone receptor positive (HR+), human epidermal growth 	<p>1. Patient shows evidence of positive response to therapy.</p> <p>2. Approval Duration: 12 months.</p>

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>factor receptor 2-negative (HER2-) advanced or metastatic breast cancer in combination with:</p> <ol style="list-style-type: none"> a. An aromatase inhibitor as initial endocrine based therapy. b. Fulvestrant in patients with disease progression following endocrine therapy. <ol style="list-style-type: none"> 2. Patient age \geq 18 years. 3. Patient has recurrent or metastatic disease; and 4. Patient has hormone receptor positive (HR+) either estrogen receptor positive and/or progesterone receptor positive disease; and 5. Patient has human epidermal growth factor receptor 2 (HER2)-negative breast cancer; AND 6. Patient meets one of the following: <ul style="list-style-type: none"> • Patient is post-menopausal; or • Patient is pre/peri-menopausal and has had either surgical bilateral oophorectomy or ovarian irradiation OR is receiving ovarian suppression/ablation with a GnRH. 7. Ibrance will be used in combination with one of the following: anastrozole, exemestance, letrozole, or fulvestrant. 8. Medication ordered by an Oncologist 9. Approval Duration: 12 months. 	
palopegteriparatide (Yorvipath) 168 mcg/0.56 ml 294 mcg/0.98 ml 420 mcg/1.4 ml	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Treatment of hypoparathyroidism in adults. 2. Patient has had hypoparathyroidism for \geq 6 months. 3. Patient has documentation or claims history supporting treatment with a vitamin D metabolite/analog therapy with calcitriol \geq 0.5 mcg per day or alfacalcidol \geq 1.0 mcg per day. 4. Patient is treated with elemental calcium at doses \geq 800 mg per day. 	<ol style="list-style-type: none"> 1. Documentation of positive clinical benefit from therapy as evidenced by the maintenance or normalization of calcium levels compared to baseline. 2. Approval Duration: 12 months.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ol style="list-style-type: none"> 5. Serum 25-hydroxyvitamin D concentration is above the lower limit of normal laboratory range. 6. Laboratory results confirming albumin-corrected serum calcium is ≥ 7.8 mg/dL prior to initiation of therapy. 7. Laboratory results confirming magnesium level is within normal laboratory limits. 8. Not prescribed for acute post-surgical hypoparathyroidism (within six months of surgery) and expected recovery from hypoparathyroidism. 9. Approval Duration: 12 months. 	
patisiran (Onpattro) Solution 10mg/5ml	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Treatment of polyneuropathy in adults with hereditary transthyretin-mediated (hATTR) amyloidosis. 2. Patient age ≥ 18 years. 3. Medication ordered by a rheumatologist, neurologist, or a specialist in the treatment of amyloidosis. 4. Diagnosis of hATTR with polyneuropathy confirmed by the presence of a transthyretin (TTR) gene mutation (e.g., V30M, A97S, T60A, E89Q, S50R). 5. Documentation of one of the following baseline tests: <ul style="list-style-type: none"> • Modified Neuropathy Impairment Scale +7 (mNIS+7) composite score. • Polyneuropathy disability (PND) score of \leq IIIb • Familial amyloid polyneuropathy (FAP) Stage 1 or 2 6. Patient has clinical signs and symptoms of polyneuropathy (i.e., weakness, sensory loss, decreased motor strength, decreased gait speed) 7. Other causes of peripheral neuropathy have been assessed and ruled out. 	<ol style="list-style-type: none"> 1. Patient continues to meet the initial approval criteria. 2. Documentation of therapeutic response as evidenced by the stabilization or improvement from baseline in one of the following: <ul style="list-style-type: none"> • mNIS+7 score • polyneuropathy disability (PND) score \leq IIIb • familial amyloid polyneuropathy (FAP) Stage 1 or 2. 3. Approval Duration: 12 months.

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	8. Patient will not be receiving Onpattro in combination with oligonucleotide agents (Onpattro, Tegsedi) 9. Prescribed by, or in consultation with, a neurologist, geneticist, or physician specializing in the treatment of amyloidosis. 10. Approval Duration: 12 months	
pegcetacoplan (Empaveli) injection 1080mg/20ml	1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Treatment of adult patients with paroxysmal nocturnal hemoglobinuria (PNH). 2. Documentation supporting diagnosis of PNH as confirmed by both of the following: <ul style="list-style-type: none"> • Flow cytometry analysis confirming presence of PNH clones; and • Laboratory results, signs and/or symptoms attributed to PNH (e.g., abdominal pain, anemia, dyspnea, extreme fatigue, smooth muscle dystonia, unexplained or unusual thrombosis, hemolysis/hemoglobinuria, kidney disease, pulmonary hypertension, etc.) 3. Patient age ≥ 18 years. 4. Patient is not receiving Empaveli in combination with another complement inhibitor used for the treatment of PNH (e.g., Soliris, Ultomiris); OR 5. Patient is currently receiving Soliris (eculizumab) which will be discontinued after an initial 4-week overlap period with Empaveli; OR 6. Patient is currently receiving Ultomiris (ravulizumab-cwvz) which will be discontinued and Empaveli will be initiated no more than 4 weeks after the last dose. 7. Prescribed by either a hematologist or oncologist. 8. Approval Duration: 6 months.	1. Documentation of positive clinical response to Empaveli therapy (e.g, increased or stabilization of hemoglobin levels, reduction in transfusions, improvement in hemolysis, decrease in LDH, increased reticulocyte count, etc.) 2. Patient is not receiving Empaveli in combination with another complement inhibitor used for the treatment of PNH (e.g., Soliris, Ultomiris). 3. Prescribed by or in consultation with an oncologist or hematologist. 4. Approval Duration: 12 months.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
pegloticase (Krystexxa) injection solution 8mg/ml	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Treatment of chronic gout in adults refractory to conventional therapy. 2. Verified there is no G6PD deficiency prior to therapy initiation. 3. Patient aged 18 years or older. 4. Not for the treatment of asymptomatic hyperuricemia. Patient as symptomatic gout. 5. Inadequate treatment response, intolerance, or contraindication to ONE of the following: allopurinol or probenecid. 6. Oral anti-hyperuricemic agents are discontinued. 7. Prescriber agrees to monitor serum uric acid levels prior to subsequent infusions and consider discontinuing treatment if levels rebound and exceed 6 mg/dl. 8. Medication ordered by Rheumatologist, Nephrologist or Podiatrist. 9. Approval duration: 3 months. 	<ol style="list-style-type: none"> 1. Patient must have chronic, symptomatic gout. 2. Documented improvement in serum uric acid level 3. NO glucose-6-phosphate dehydrogenase (G6PD) deficiency 4. Renewal request may be denied if patient has 2 or more consecutive uric acid levels above 6 mg/dl. 5. Renewal duration: 3 months.
ponatinib (Iclusig) tablets 10mg, 15mg, 30mg, 45mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Chronic phase (CP) chronic myeloid leukemia (CML) with resistance or intolerance to at least 2 prior kinase inhibitors. • Accelerated phase (AP) or blast phase (BP) CML or Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ ALL) for whom no other kinase inhibitors are indicated. • T315I-positive CML (chronic-, accelerated-, or blast phase) or T315I-positive Ph+ ALL. 2. Medication ordered by an Oncologist. <p>Acute Lymphoblastic Leukemia:</p>	<ol style="list-style-type: none"> 1. Patient shows positive clinical response to therapy. 2. Patient has not experienced any severe adverse effects from therapy. 3. Approval Duration: 12 months.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ol style="list-style-type: none"> 1. Patient is ≥ 15 years of age; AND 2. Patient has Philadelphia chromosome-positive acute lymphoblastic leukemia; AND 3. Patient meets ONE of the following: <ul style="list-style-type: none"> • The drug will be used in combination with chemotherapy; or • The acute lymphoblastic leukemia is T315I-positive; OR • The patient has tried at least one other tyrosine kinase inhibitor that is used for Ph+ ALL (e.g., Sprycel (dasatinib)). 4. Approval Duration: 12 months. <p><u>Chronic Myeloid Leukemia (CML):</u></p> <ol style="list-style-type: none"> 1. Patient is ≥ 18 years; AND 2. Patient has Philadelphia chromosome-positive chronic myeloid leukemia; AND 3. Patient meets ONE of the following: <ul style="list-style-type: none"> • The chronic myeloid leukemia is T315I-positive; OR • Patient has tried at least two other tyrosine kinase inhibitors indicated for use in Ph+ CML (e.g., imatinib, dasatinib, nilotinib); OR • Patient meets BOTH of the following: <ul style="list-style-type: none"> ○ Patient has accelerated-phase CML or blast-phase CML; AND ○ No other tyrosine kinase inhibitor is indicated. 4. Approval Duration: 12 months. <p>COVERED COMPENDIAL USES: <u>Gastrointestinal Stromal Tumor:</u></p> <ol style="list-style-type: none"> 1. Patient age ≥ 18 years; and 	

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>2. Patient has tried each of the following four therapies:</p> <ul style="list-style-type: none"> • One of either imatinib or avapritinib; AND • One of either sunitinib or dasatinib; AND • Stivarga (regorafenib); AND • Qinlock (repreтинib). <p>3. Approval Duration: 1 year.</p> <p><u>Myeloid/Lymphoid Neoplasms with Eosinophilia:</u></p> <p>1. Patient age ≥ 18 years; and</p> <p>2. Patient meets ONE of the following:</p> <ul style="list-style-type: none"> • The tumor has an ABL1 rearrangement, OR • The tumor has an FGFR1 rearrangement. <p>3. Approval Duration: 12 months.</p>	
<p>posaconazole (Noxafil) 40mg/ml suspension</p> <p>100 mg tablets</p>	<p>1. Ordered for an approved indication for use:</p> <ul style="list-style-type: none"> • Treatment of invasive aspergillosis in adults and pediatric patients ≥ 13 years of age. (Injection and tablets). • Treatment of oropharyngeal candidiasis (OPC), including OPC refractory (rOPC) to itraconazole and/or fluconazole in adults or pediatric patients ≥ 13 years of age. • Prophylaxis of invasive Aspergillus and Candida infections in patients at high risk of infection development due to being severely <p>3. The patient is being prescribed for the treatment for the prevention of invasive Aspergillus and Candida infections in a patient who is at high risk of developing these infections due to being severely immunocompromised; OR</p> <p>4. The patient is being prescribed injection or delayed-release tablets for the treatment of invasive aspergillosis; OR</p> <p>5. The patient is being prescribed oral suspension for</p>	<ul style="list-style-type: none"> • Patient at high infection risk: severely immunocompromised, such as HSCT recipients with GVHD or those with hematologic malignancies with prolonged neutropenia from chemotherapy. • Oral suspension is not substitutable with tablets or PowderMix oral suspension due to differences in dosing of each formulation. • Coadministration is Contraindicated with sirolimus, ergot alkaloids, HMG-CoA reductase inhibitors. • Significant risk for drug-drug interactions.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>the treatment of moderate to severe oropharyngeal candidiasis AND:</p> <ul style="list-style-type: none"> • The patient has experienced an inadequate treatment response to fluconazole, OR • The patient has experienced an intolerance to fluconazole, OR • The patient has a contraindication that would prohibit a trial of fluconazole. <p>6. Approval Duration: up to 12 months.</p>	
<p>ravulizumab-cwvz (Ultomiris) injection solution 300mg/mL, 1100mg/11ml</p>	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • treatment of adult and pediatric patients \geq 1 month of age with paroxysmal nocturnal hemoglobinuria (PNH). • treatment of adult and pediatric patients \geq 1 month of age with atypical hemolytic uremic syndrome (aHUS) to inhibit complement-mediated thrombotic microangiopathy (TMA). • treatment of adult patients with generalized myasthenia gravis (gMG) who are anti-acetylcholine receptor (AChR+) antibody positive. 2. FDA approved patient age. 3. Documentation to support diagnosis: <p>PNH:</p> <ul style="list-style-type: none"> • Flow cytometric confirmation of PNH type III red cells; AND • Patient had at least one transfusion in the preceding 24 months; OR • Documented history of major adverse thrombotic vascular events from thromboembolism; OR • Patient has high disease activity defined as lactic dehydrogenase (LDH) level \geq 1.5 times the upper limit of normal with one of the following 	<ol style="list-style-type: none"> 1. Clinical documentation must be provided to confirm that current criteria are met and that the medication is providing clinical benefit. 2. gMG: Improvement and maintenance of at least a 2-point improvement (reduction) in the MG-ADL score from pre-treatment baseline and reduction of signs and symptoms of MG required to show clinical benefit. 3. NOTE: dose escalation of immunosuppressive therapy, or additional rescue therapy from baseline to treat MG or exacerbation of symptoms during use is considered treatment failure. 4. Not receiving in combination with Empaveli or Soliris. 5. Approval Duration: up to 12 months.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>symptoms: weakness, fatigue, hemoglobinuria, abdominal pain, dyspnea, hemoglobin, 10 g/dL, a major vascular event, dysphagia, or erectile dysfunction.</p> <ul style="list-style-type: none"> • History of failure to/contraindication or intolerance to Empaveli therapy; • Patient age < 18 years or currently pregnant. <p>aHUS:</p> <ul style="list-style-type: none"> • Common causes of aHUS have been ruled out, including infectious causes of HUS and thrombotic thrombocytopenic purpura (TTP). • Ultomiris is not indicated for the treatment of patients with Shiga toxin E. coli related hemolytic uremic syndrome (STEC-HUS). • Must present with the following symptoms: <ul style="list-style-type: none"> ○ Hemoglobin < 10 g/dL ○ Platelets, 150,000/mm³ ○ Documented evidence of hemolysis, such as elevated LDH levels, decreased haptoglobin level or schistocytosis. • Increased serum creatinine OR currently undergoing dialysis. <p>gMG:</p> <ul style="list-style-type: none"> • Patient has not failed previous course of Soliris or Ultomiris therapy; • Positive serologic test for anti-AChR antibodies; AND • One of the following: <ul style="list-style-type: none"> ○ History of abnormal neuromuscular transmission test demonstrated by single-fiber electromyography (SFEMG) or repetitive nerve stimulation OR 	

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> ○ History of positive anticholinesterase test (e.g. edrophonium chloride test) OR ○ Pt has demonstrated improvement in MG signs on oral cholinesterase inhibitors as assessed by the treating neurologist; AND ● Patient has MGFA clinical classification of II, III, or IV at initiation of treatment; AND ● Patient has Myasthenia gravis-specific activities of daily living scale (MG-ADL) total score ≥ 6 at initiation of treatment; AND ● One of the following: <ul style="list-style-type: none"> ○ History of failure to at least two immunosuppressive agents over the previous 12-months (e.g., azathioprine, mtx, cyclosporin, mycophenolate); OR ○ History of failure to at least one immunosuppressive therapy and has required four or more courses of plasmapheresis/plasma exchanges, and/or intravenous immune globulin over the previous 12 months without symptom control; AND <p>4. Recommended vaccinations at least 2 weeks prior to administration of first dose Ultomiris.</p> <p>5. Cannot be used in combination with other medications in the same class, such as Soliris.</p> <p>6. Medication ordered by Hematologist, Nephrologist, or Oncologist registered with Ultomiris REMS program.</p> <p>7. Approval Duration: 12 months.</p>	
resmetirom (Rezdiffra) tablets 80 mg, 100 mg	1. Ordered for an approved indication for use:	1. Patient meets ONE of the following: <ul style="list-style-type: none"> ● Completed ≥ 1 year and < 2 years of therapy with Rezdiffra AND the patient

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
(60 mg is non-formulary)	<ul style="list-style-type: none"> • Treatment of adults with nonalcoholic steatohepatitis (NASH/MASH) with moderate to advanced (F2 or F3) liver fibrosis. <ol style="list-style-type: none"> 2. Patient age ≥ 18 years, AND 3. Prior to treatment, the diagnosis of MASH/NASH is confirmed by one of the following: <ul style="list-style-type: none"> • Patient has had a liver biopsy AND meets both of the following: <ul style="list-style-type: none"> ○ Liver biopsy was performed within the 6 months preceding treatment with Rezdifra; AND ○ Liver biopsy shows non-alcoholic fatty liver disease activity score ≥ 4 with a score > 1 in ALL of the following: steatosis, ballooning, and lobular inflammation OR • Patient has had ONE of the following imaging exams performed within the 3 months preceding treatment with Rezdifra: <ul style="list-style-type: none"> ○ Elastography (e.g. Fibroscan, transient elastography, magnetic resonance elastography, acoustic radiation force impulse imaging, or shear wave elastography); OR ○ Computed tomography; OR ○ Magnetic resonance imaging. 4. Patient meets ONE of the following prior to treatment with Rezdifra: <ul style="list-style-type: none"> • Patient has Stage F2 fibrosis; OR • Patient has Stage F3 fibrosis; AND THREE or more of the following metabolic risk factors that are managed according to Standards of Care: <ul style="list-style-type: none"> ○ Central obesity 	<p>has derived benefit from treatment as demonstrated by at least ONE of the following: MASH/NASH resolution AND no worsening of fibrosis OR</p> <ul style="list-style-type: none"> • No worsening of MASH/NASH AND improvement in fibrosis by ≥ 1 stage; OR • Patient has completed ≥ 2 years of treatment AND the patient has not had worsening of fibrosis or MASH/NASH AND according to the prescriber, the patient has not progressed to stage F4 (cirrhosis). <ol style="list-style-type: none"> 2. Metabolic risk factors are managed according to standard of care; AND 3. According to the prescriber, the patient meets ONE of the following: <ul style="list-style-type: none"> • <u>Female patients</u>: Alcohol consumption < 20 grams per day; OR • <u>Male patients</u>: Alcohol consumption < 30 grams per day. <p><i>Note: One standard drink (or one alcoholic drink equivalent) contains ~14 grams of pure alcohol, which is found in 12 ounces of regular beer, 5 ounces of wine, or 1.5 ounces of distilled spirits.</i></p> <ol style="list-style-type: none"> 4. This indication is approved under accelerated approval based on overall response rate and duration of response. Continued approval for this indication may be contingent upon

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> ○ Hypertriglyceridemia ○ Reduced high-density lipoprotein cholesterol, ○ Hypertension ○ Elevated fasting plasma glucose indicative of diabetes or pre-diabetes; AND <p>5. According to the prescriber, the patient meets ONE of the following:</p> <ul style="list-style-type: none"> • <u>Female patients</u>: Alcohol consumption < 20 grams per day; OR • <u>Male patients</u>: Alcohol consumption < 30 grams per day. <p><i>Note: One standard drink (or one alcoholic drink equivalent) contains ~14 grams of pure alcohol, which is found in 12 ounces of regular beer, 5 ounces of wine, or 1.5 ounces of distilled spirits.</i></p> <p>6. Other causes of liver disease or hepatic steatosis have been ruled out (e.g., alcoholic steatohepatitis, acute fatty liver, autoimmune hepatitis, Hepatitis A, B, or C, hemochromatosis, drug-induced liver disease, etc.), AND</p> <p>7. Provider attestation that member has adopted liver-protective lifestyle interventions such as optimizing weight loss, dietary changes, and exercise, AND</p> <p>8. Member does not have evidence of cirrhosis, hepatic decompensation, or hepatocellular carcinoma (HCC).</p> <p>9. All other indications are excluded from coverage as experimental.</p> <p>10. Prescribed by, or in consultation with an endocrinologist, hepatologist or gastroenterologist.</p>	<p>verification and description of clinical benefit in confirmatory trials.</p> <p>5. Approval Duration: 12 months.</p>

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
riociguat (Adempas) 0.5 mg, 1 mg, 1.5 mg, 2 mg, 2.5 mg tablets	11. Approval Duration: 12 months 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Persistent/recurrent Chronic Thromboembolic Pulmonary Hypertension (CTEPH) (WHO Group 4) after surgical treatment or inoperable CTEPH to improve exercise capacity and WHO functional class. • Pulmonary Arterial Hypertension (PAH) (WHO Group 1) to improve exercise capacity, improve WHO functional class and to delay clinical worsening. 2. Will not be used in combination with a phosphodiesterase 5 inhibitor (PDE5I; e.g. sildenafil, tadalafil) CTEPH: <ol style="list-style-type: none"> 1. Patient is diagnosed with inoperable or persistent/recurrent chronic thromboembolic pulmonary hypertension (CTEPH); AND 2. CTEPH is symptomatic; AND 3. Prescribed by or in consultation with a cardiologist, pulmonologist, or rheumatologist. 4. Approval Duration: 12 months. PAH: <ol style="list-style-type: none"> 1. Patient has symptomatic PAH. 2. Diagnosis of PAH is confirmed by right heart catheterization. 3. For patients with WHO functional class I PAH: patient has previous trials of at least one PDE5I medication (sildenafil, tadalafil) AND at least one endothelin receptor agonist medication (ambrisentan, macetentan, bosentan) 4. For patients with WHO functional class II-IV PAH: 	<ol style="list-style-type: none"> 1. Clinical documentation supports that the patient is receiving clinical benefit from Adempas therapy. 2. Approval Duration: 12 months.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>patient has previously tried and failed OR has a contraindication to using a PDE5I medication, as part of combination oral therapy with one or more PAH medications from other classes.</p> <p>5. Prescribed by or in consultation with a cardiologist, pulmonologist, or rheumatologist.</p> <p>6. Approval Duration: 12 months.</p>	
<p>ruxolitinib (Jakafi) tablets 5mg, 10mg, 15mg, 20mg, 25mg</p>	<ol style="list-style-type: none"> Ordered for an approved indication for use: <ul style="list-style-type: none"> Intermediate or high-risk myelofibrosis, including primary myelofibrosis, post-polycythemia vera myelofibrosis and post-essential thrombocythemia myelofibrosis in adults. Polycythemia vera in adults who have had an inadequate response to or are intolerant of hydroxyurea. Steroid-refractory acute graft-versus-host disease in adult and pediatric patients 12 years and older. Chronic graft-versus-host disease after failure of one or two lines of systemic therapy in adult and pediatric patients 12 years and older. Medication ordered by Hematologist or Oncologist. Approval Duration: 12 months. 	<p>Limitations of Use:</p> <ul style="list-style-type: none"> Avoid concomitant use with fluconazole doses greater than 200 mg. Reduce Jakafi dosage with fluconazole doses ≤ 200 mg. Strong CYP3A4 Inhibitors: Reduce, interrupt, or discontinue Jakafi doses as recommended except in patients with acute or chronic graft-versus-host-disease.
<p>sastralizumab-mwge (Enspryng) injection solution 120mg/ml</p>	<ol style="list-style-type: none"> Ordered for an approved indication for use: <ul style="list-style-type: none"> Treatment of neuromyelitis optica spectrum disorder (NMOSD) in adult patients who are anti-aquaporin-4 (AQP4) antibody positive. Must submit FDA-approved testing showing antibody positive. Age ≥ 18 years. Prescriber attests that baseline evaluation has been done and there are no contraindications to use (e.g., Hep B, TB, LFT's, live or live-attenuated vaccines 4 	<ol style="list-style-type: none"> Meets all initial criteria, AND Provider attestation of continued benefit. Use in caution if ALT/AST > 1.5 x ULN. Contraindicated in patients with active hepatitis B infection or active or untreated latent tuberculosis. Approval duration: 12 months

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>weeks prior or 2 weeks for non-live vaccines).</p> <ol style="list-style-type: none"> 5. Prescriber attests that subsequent appropriate evaluation and monitoring will be done based on the package insert (e.g., infections, LFT's, CBCs – neutrophils) 6. Medication ordered by neurologist, immunologist, or ophthalmologist experienced in treatment of this disease. 7. Approval Duration: 12 months. 	
<p>secukinumab (Cosentyx)</p> <p>75 mg SOSY, 150 mg SOSY, Sensoready 150 mg pens Unoready 300 mg pens</p>	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use following the indication-specific criteria as outlined below. Please note that the following indications are NOT approved for coverage: Crohn's Disease, Rheumatoid Arthritis, or Uveitis. 2. May not be ordered for concurrent use with a biologic or targeted synthetic oral small molecule drug (e.g., TNF inhibitors, Inhibitors of interleukin types 1, 6, 12, 17, 17A, 23, or combinations thereof, CD20-directed cytolytic antibodies, JAKs, PDE4s, Sphingosine 1 phosphate receptor modulators due to increased risk of adverse effects and lack of clinical data supporting additive efficacy. <p>Ankylosing Spondylitis:</p> <ul style="list-style-type: none"> • Patient age ≥ 18 years; AND • Prescribed by or in consultation with a rheumatologist • Approval Duration: 6 months. <p>Enthesitis-Related Arthritis:</p> <ul style="list-style-type: none"> • Patient age ≥ 4 years of age; AND • Prescribed by or in consultation with a rheumatologist • Approval Duration: 6 months. 	<p>Ankylosing Spondylitis:</p> <ul style="list-style-type: none"> • Patient has been established on Cosentyx SQ or IV for at least 6 months; AND • Patient shows positive clinical response by way of at least one objective measure or improvement in at least one symptom. • Approval Duration: 12 months. <p>Enthesitis-Related Arthritis:</p> <ul style="list-style-type: none"> • Patient has been established on Cosentyx SQ for at least 6 months; AND • Patient shows positive clinical response by way of at least one objective measure; AND • Patient shows positive clinical response or improvement in at least one symptom. • Approval Duration: 12 months. <p>Hidradenitis Suppurativa:</p>

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p><u>Hidradenitis Suppurativa:</u></p> <ul style="list-style-type: none"> • Patient age ≥ 18 years; AND • Patient has tried at least one other therapy (e.g. corticosteroids, systemic antibiotics, or isotretinoin). • Prescribed by or in consultation with a dermatologist. • Approval Duration: 3 months. <p><u>Non-Radiographic Axial Spondylarthritis:</u></p> <ul style="list-style-type: none"> • Patient age ≥ 18 years; AND • Patient has objective signs of inflammation, defined as at least ONE of the following: <ul style="list-style-type: none"> ○ C-reactive protein elevated beyond the upper limit of normal for the reporting laboratory, OR ○ Sacroiliitis reported on magnetic resonance imaging; AND • Prescribed by or in consultation with a rheumatologist • Approval Duration: 6 months. <p><u>Plaque Psoriasis:</u></p> <ul style="list-style-type: none"> • Patient age ≥ 6 years; AND • Patient meets ONE of the following conditions: <ul style="list-style-type: none"> ○ Patient has tried at least one traditional systemic agent for psoriasis for at least 3 months, unless intolerant (e.g. methotrexate, cyclosporine, or acitretin or a 3-month trial of psoralen plus ultraviolet light unless patient has already had a 3-month trial or previous intolerance to at least one biologic other than Cosentyx/biosimilar); OR 	<ul style="list-style-type: none"> • Patient has been established on Cosentyx SQ for at least 3 months; and • Patient has experienced positive clinical response to at least one objective measure from baseline (e.g. Hurley staging, Sartorius score, Physician Global Assessment, and Hidradenitis Suppurativa Severity index); AND • Patient has experienced positive clinical response in at least one symptom (e.g. decreased pain or drainage of lesions, nodules, or cysts). • Approval Duration: 12 months. <p><u>Non-Radiographic Axial Spondylarthritis:</u></p> <ul style="list-style-type: none"> • Patient has been established on Cosentyx SQ or IV for at least 6 months; AND • Patient shows positive clinical response by way of at least one objective measure or improvement in at least one symptom. • Approval Duration: 12 months. <p><u>Plaque Psoriasis:</u></p> <ul style="list-style-type: none"> • Patient has been established on Cosentyx SQ for at least 3 months. • Patient has experienced a positive clinical response defined as improvement from baseline in at

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> ○ Patient has a contraindication to methotrexate ● Prescribed by or in consultation with a dermatologist. ● Approval Duration: 3 months. <p>Psoriatic Arthritis:</p> <ul style="list-style-type: none"> ● Patient age ≥ 2 years; AND ● Prescribed by or in consultation with a rheumatologist or dermatologist. ● Approval Duration: 6 months. 	<p>least one of the following: estimated affected BSA, erythema, induration/thickness and/or scale of areas affected by psoriasis.</p> <ul style="list-style-type: none"> ● Patient has experienced a positive clinical response in at least one symptom such as decreased pain, itching, and/or burning. ● Approval Duration: 12 months. <p>Psoriatic Arthritis:</p> <ul style="list-style-type: none"> ● Patient has been established on Cosentyx SQ or IV for at least 6 months; AND ● Patient shows positive clinical response by way of at least one objective measure or improvement in at least one symptom. ● Approval Duration: 12 months.
selexipag (Uptravi) tablets 200 mcg, 400 mcg, 600 mcg, 800 mcg, 1000 mcg, 1200 mcg, 1400 mcg, 1600 mcg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> ● Treatment of pulmonary arterial hypertension (PAH) (WHO Group I) to delay disease progression and reduce the risk for hospitalization for PAH. 2. Patient aged ≥ 18 years. 3. Patient diagnosed with pulmonary hypertension WHO group 1. 4. Patient has had a right heart catheterization and the diagnosis of WHO Group 1 PAH is confirmed. 5. Patient meets one of the following criteria (a or b): <ol style="list-style-type: none"> a. Patient has tried or is currently receiving at least one oral medication for PAH from one of the three following different categories (either alone or in combination) each for ≥ 60 days: one 	<ol style="list-style-type: none"> 2. Patient meets initial approval criteria. 3. Approval Duration: 12 months.

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>phosphodiesterase type 5 (PDE5) inhibitor (i.e. sildenafil or tadalafil), one endothelin receptor antagonist (ERA) (i.e., bosentan, ambrisentan or macitentan), or Adempas (riociguat) OR</p> <p>b. Patient is currently receiving, or has a history of receiving, one prostacyclin therapy for PAH (i.e., Tyvaso or Orenitram, (Treprostinil), Ventavis (iloprost), or epoprostenol).</p> <p>6. May not concurrently be prescribed Orenitram, inhaled prostacyclin products, or parenteral prostacyclin agents used for PAH (e.g. Tyvaso, Ventavis, epoprostenol, Treprostinil SQ or IV [Remodulin, generics]).</p> <p>7. May not have Child-Pugh Class C or D liver disease.</p> <p>8. May not be on dialysis or have eGFR < 15 ml/min</p> <p>9. Prescribed by or in consultation with a cardiologist or pulmonologist.</p> <p>10. Quantity Limits: 1 titration/starter pack per 365 days Max 2 tablets per day and total daily dose of 3200 mcg</p> <p>11. Approval Duration: 12 months.</p>	
selpercatinib (Retevmo) capsules 40mg, 80mg	1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Adult patients with locally advanced or metastatic non-small cell lung cancer (NSCLC) with a rearranged during transfection (RET) gene fusion, as detected by an FDA- approved test. • Adult and pediatric patients ≥ 12 years of age with advanced or metastatic medullary thyroid cancer (MTC) with a RET mutation, who require systemic therapy. • adult and pediatric patients ≥ 12 years of age with advanced or metastatic RET fusion-positive 	4. Patient does not show evidence of progressive disease while on Retevmo therapy. 5. Approval Duration: 12 months.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>thyroid cancer who require systemic therapy and who are radioactive iodine-refractory (if radioactive iodine is appropriate).</p> <ul style="list-style-type: none"> • Adult patients with locally advanced or metastatic solid tumors with a RET gene fusion that have progressed on or following prior systemic treatment or who have no satisfactory alternative treatment options. <p>2. Medication ordered by an Oncologist.</p> <p>3. Approval Duration: 12 months.</p>	
<p>semaglutide (Ozempic, Rybelsus)</p> <p>Ozempic 2mg/3 ml (0.25 mg or 0.5 mg/week; 4mg/3ml (1 mg per week) 8mg/3ml (2 mg per week)</p> <p>Rybelsus 3mg, 7mg, 14mg</p>	<p>1. Ordered for an approved indication for use:</p> <ul style="list-style-type: none"> • As adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes mellitus. • Treatment of adult patients with Type 2 diabetes mellitus for risk reduction of major cardiovascular events (cardiovascular death, nonfatal myocardial infarction, nonfatal stroke) who have established CVD or multiple cardiovascular risk factors. <p>2. Patient age \geq 18 years. NOTE: <i>Semaglutide is approved in adolescents for weight loss only and is not a covered benefit.</i></p> <p>3. Patient must have Type 2 Diabetes Mellitus ***NOTE: <i>Type 1 DM does NOT qualify for coverage***</i></p> <p>4. A1c or CGM Time in Range% (TIR%) report within previous 3 months.</p> <p>5. A urine albumin-to-creatinine ratio (uACR) within the previous 12 months.</p> <p><u>Treatment of Type 2 Diabetes without regard to CVD risk factors:</u></p>	<p>Cannot be approved for indication of weight management.</p> <p>1. Rybelsus 3 mg dose may not be renewed and must be escalated to 7 mg or 14 mg dose.</p> <p>2. Chart notes with A1c or CGM report with TIR% within previous 3 months.</p> <p>3. A urine albumin-to-creatinine ratio (uACR) within the previous 12 months.</p> <p>2. Documented positive clinical response defined as one of the following:</p> <p><u>Baseline (pre-GLP-1) A1c was \geq 8.0 and:</u></p> <ul style="list-style-type: none"> • A1c has decreased by \geq 1% since onset of therapy or • TIR% was \leq 55% and has increased \geq 10% <p><u>Baseline (pre-GLP-1) A1c was \geq 6.5 but $<$ 8.0 and:</u></p> <ul style="list-style-type: none"> • A1c or TIR% has improved. • Not eligible for renewal if A1c has not changed or has increased or

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>The patient has an A1c (hemoglobin A1c) of ≥ 7.5 (TIR $\leq 60\%$).</p> <p style="text-align: center;">OR</p> <p><u>Treatment of Type 2 Diabetes with CVD as defined below:</u></p> <ul style="list-style-type: none"> • Pre-treatment A1c is ≥ 6.5 (TIR $\leq 70\%$) AND • BMI ≥ 27 kg/m² (documentation within previous 90 days of current height and weight); AND <p>Documentation submitted to show that the patient has at least ONE of the following:</p> <ul style="list-style-type: none"> • Prior myocardial infarction; OR • Prior stroke (ischemic or hemorrhagic); OR • Symptomatic peripheral arterial disease (PAD) as evidenced by: <ul style="list-style-type: none"> ○ Intermittent claudication with ankle-brachial index (ABI) less than 0.85 (at rest); OR ○ Peripheral arterial revascularization procedure; OR ○ Amputation due to atherosclerotic disease. <p>7. May not be concurrently using:</p> <ul style="list-style-type: none"> • ANY other GLP1 or GLP1/GIP combination drug (e.g., Mounjaro, Ozempic, Rybelsus, Saxenda, Soliqua, Trulicity, Victoza (liraglutide), Wegovy, Xultrophy or Zepbound). • ANY DPP4i (e.g., alogliptin, Januvia (sitagliptin), Tradjenta (Linagliptin), Onglyza (saxagliptin)). • Agents for severe constipation: metoclopramide, Amitiza (lubiprostone), Linzess (linaclotide), Motegrity (prucalopride) or Trulance (plecanatide). 	<p>TIR% has decreased.</p> <p>3. Patient has not had medical intervention for:</p> <ul style="list-style-type: none"> • Pancreatitis, or • Severe gastrointestinal events (e.g., hospitalization or new start GI motility agent). <p>4. May not be concurrently using:</p> <ul style="list-style-type: none"> • ANY other GLP1 or GLP1/GIP combination drug (e.g., liraglutide, Ozempic, Rybelsus, Saxenda, Soliqua, Trulicity, Victoza, Xultrophy or Zepbound) AND/OR • ANY DPP4i (e.g., alogliptin, Januvia (sitagliptin), Onglyza (saxagliptin), or Tradjenta (linagliptin)). • Agents for severe constipation: metoclopramide, Amitiza (lubiprostone), Linzess (linaclotide), Motegrity (prucalopride) or Trulance (plecanatide). <p>8. PBM claims data shows consistent adherence as shown by no instance of a drug-free interval greater than 2 months at which time the patient would need to satisfy the initial criteria.</p> <p>9. Approval Duration: 12 months</p>

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>8. Prescriber attests that medication is prescribed in accordance with prescribing information, including screening for any black box warnings and all contraindications.</p> <p>9. May not be approved for patients with:</p> <ul style="list-style-type: none"> • Personal or family history of medullary thyroid carcinoma (MTC) or in patients with Multiple Endocrine Neoplasia syndrome type 2 (MEN 2). • Current pregnancy; and/or • A history of confirmed pancreatitis, and/or • Suicidal ideation or new onset depression. <p>10. Starter doses are limited and require dose escalation. Starter doses are defined as:</p> <ul style="list-style-type: none"> • Ozempic: the 0.25/0.5 mg strength combines the starter-dose and titration-dose and is limited to two, 28-day dispenses and then must be dose escalated to 1 mg per week dose UNLESS A1c ≤ 7.0 or TIR ≥ 65%. • Rybelsus 3 mg is a starter dose and limited to one, 30-day dispense then must be dose escalated to 7 mg. <p>11. Ozempic: Limited to a maximum of 1 pen per 28 days. Rybelsus: Limited to 30 capsules per 30 days.</p> <p>12. Maximum approval duration: 12 months.</p>	
<p>sildenafil (Revatio) 20 mg tablets 10 mg/ml solution</p> <p><i>**NOTE: sildenafil 25 mg, 50 mg and 100 mg dosage forms are indicated for erectile dysfunction <u>only</u> and are not covered by the formulary.</i></p>	<p>1. Ordered for an approved indication for use:</p> <ul style="list-style-type: none"> • treatment of pulmonary arterial hypertension (PAH) (WHO Group I) in adults to improve exercise ability and delay clinical worsening. <p>2. Patient is not concurrently on organic nitrates (i.e. isosorbide mononitrate, isosorbide dinitrate, nitroglycerin) or Adempas (riociguat), OR tadalafil, AND</p>	<p>1. All initial criteria continue to be met.</p> <p>2. Has documented positive clinical response to sildenafil treatment as determined by one or more of the following:</p> <ul style="list-style-type: none"> • Progress towards improvement in WHO functional class status,

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ol style="list-style-type: none"> 3. The diagnosis of PAH is documented by right-heart catheterization with ALL of the following: <ul style="list-style-type: none"> • Mean pulmonary artery pressure (mPAP) > 20 mmHg, • Pulmonary arterial wedge pressure (PAWP) ≤ 15 mmHg; AND • Pulmonary vascular resistance (PVR) ≥ 3 wood units. 4. Prior Authorization not required for solution for children less than 6 years of age. Tablets are preferred dosage form; solution should only be utilized when tablets cannot satisfy medical necessity. 5. May not be approved for the treatment of erectile dysfunction (ED). 6. Medication ordered by a cardiologist or pulmonologist. 7. Total daily dosage does not exceed 60 mg. 8. Approval duration: 12 months. 	<ul style="list-style-type: none"> • Improvement in right-ventricular function (based on echocardiogram or cardiac MRI), • Improvement from baseline on the 6-minute walk distance (6MWD), • Improvement in B-type natriuretic peptide plasma levels (NT-proBNP) <ol style="list-style-type: none"> 3. Approval Duration: 12 months.
solriamfetol (Sunos) tablets 75 mg, 150 mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • narcolepsy • obstructive sleep apnea 2. If ordered for narcolepsy: <ul style="list-style-type: none"> • narcolepsy diagnosis must be confirmed by sleep study, OR provider justification is included confirming that a sleep study is not feasible, AND • must have failed at least one formulary stimulant treatment (e.g. modafinil, armodafinil) or have an intolerance or contraindication for use, AND • must be ordered to manage symptoms of excessive daytime sleepiness associated with narcolepsy, and is not being ordered to manage 	Renewal Criteria: <ol style="list-style-type: none"> 1. Initial approval criteria met. 2. Provider confirmation that patient experienced positive clinical benefit with the medication. <ul style="list-style-type: none"> • Narcolepsy – reduction in symptoms of excessive daytime sleepiness from baseline • OSA – reduction in symptoms of excessive daytime sleepiness from baseline, AND patient continues to be compliant with other treatments 3. Dose has been titrated appropriately,

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>cataplexy symptoms</p> <ol style="list-style-type: none"> 3. If ordered for obstructive sleep apnea: <ul style="list-style-type: none"> • OSA diagnosis must be confirmed by sleep study, OR provider justification is included confirming that a sleep study is not feasible, AND • Standard treatments for the underlying airway obstruction have been used for ≥1 month (CPAP, BiPAP), AND • Patient is fully compliant with their ongoing treatment(s) for the underlying airway obstruction, as confirmed by provider attestation, AND • Must have failed at least one formulary stimulant treatment (e.g. modafinil, armodafinil) or have an intolerance or contraindication for use. 4. For patients with a known past medical history of chronic kidney disease, baseline labs with eGFR from within the last 2 months submitted with the request. <ul style="list-style-type: none"> • Requested dose is appropriate for treatment initiation based on the patient’s level of kidney impairment. 5. Target maximum daily dose cannot exceed 150 mg/day. 6. Initial approval for six months. 	<p>or patient has sufficient symptom control on requested dose.</p> <ol style="list-style-type: none"> 4. Approval Duration: 12 months
<p>somatrogon (Ngenla) solution pen-injector 24mg/1.2ml; 60mg/1.2ml</p>	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Treatment of growth failure in children due to inadequate secretion of endogenous growth hormone (GH) 2. Age 3-<18 years 3. Medication ordered by or in consultation with an Endocrinologist 4. <u>Initial approval:</u> 	<ol style="list-style-type: none"> 1. Confirmation of open epiphysial growth plates as above, OR the patient has not completed prepubertal growth 2. Patient meets at least one of the following: <ul style="list-style-type: none"> • Has an annual growth velocity of at least 2 cm during most recent approval year;

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> • Confirmation of open epiphysial growth plates • Patient meets at least one of the following: <ul style="list-style-type: none"> ○ Height is at least TWO standard deviations (SD) below the mean height for normal children of same age and gender; ○ Height velocity less than 25th percentile for age. <p>5. Approval duration: 12 months</p>	<ul style="list-style-type: none"> • Is near the terminal phase of puberty and has an annual growth velocity of at least 1 cm during the most recent approval year. <p>3. Approval duration: 12 months</p> <p>Limitations of Use:</p> <ul style="list-style-type: none"> • Ngenla will not be approved for idiopathic short stature (ISS), athletic enhancement, central precocious puberty, congenital adrenal hyperplasia, constitutional delay of growth and puberty, or anti-aging purposes.
<p>somatropin [recombinant human growth hormone] (Norditropin FlexPro; Serostim) injection</p> <p>Norditropin 5/1.5ml, 10/1.5ml, 15/1.5ml, 30mg/3ml</p> <p>Serostim 4mg, 5mg, 6mg</p>	<p>1. Ordered for an approved indication for use:</p> <ul style="list-style-type: none"> • Growth failure in pediatric patients: <ul style="list-style-type: none"> ○ Due to inadequate endogenous growth hormone secretion; short stature associated with Turner Syndrome [Norditropin FlexPro] ○ Idiopathic Short Stature (ISS); short stature born small for gestational age (SGA) with no catch-up growth by age 2 to 4 years; Prader-Willi syndrome; short stature associated with Noonan syndrome[Norditropin ONLY] ○ Growth failure associated with chronic kidney disease until time of renal transplant. • Growth hormone deficiency in adults: replacement of endogenous growth hormone in adults with growth hormone deficiency 	<p>Growth failure in pediatric patients:</p> <ol style="list-style-type: none"> 1. Confirmation of open epiphysial growth plates as above, OR the patient has not completed prepubertal growth 2. Patient meets at least one of the following: <ul style="list-style-type: none"> • Has an annual growth velocity of at least 2 cm during most recent approval year; • Is near the terminal phase of puberty and has an annual growth velocity of at least 1 cm during the most recent approval year. <p>Adult indications for use:</p> <ol style="list-style-type: none"> 1. Clinical documentation indicating positive clinical response during previous 12 months <p>All ages: Approval duration: 12 months</p>

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>[Norditropin FlexPro]</p> <ul style="list-style-type: none"> Treatment of HIV patients with wasting or cachexia to increase lean body mass and body weight and improve physical endurance [Serostim ONLY] <ol style="list-style-type: none"> Medication ordered by an Endocrinologist or Infectious disease specialist (Serostim ONLY). For pediatric patients with growth failure: Confirmation of open epiphysial growth plates. Approval duration: 12 months 	
<p>tacrolimus extended-release (Envarsus XR) tablets 0.75mg, 1mg, 4mg</p>	<ol style="list-style-type: none"> Ordered for an approved indication for use: <ul style="list-style-type: none"> prophylaxis of organ rejection in kidney transplant in adult patients converted from tacrolimus immediate-release formulations in combination with other immunosuppressants. Documented evidence that the patient is unable to achieve or maintain an appropriate therapeutic drug level with immediate-release tacrolimus---Lab values must be submitted. Envarsus XR will be used in combination with other immunosuppressant medications to prevent kidney transplant rejection. Patient has not been diagnosed with congenital long Qt-syndrome. Prescribed by a Nephrologist and Transplant Specialist. Approval Duration: 12 months 	<ol style="list-style-type: none"> Patient has continued care with a nephrologist or transplant specialist. Patient continues to meet the initial approval criteria. No clinical evidence of organ failure. Individual has not developed any significant adverse drug effects that may exclude continued use such as: <ul style="list-style-type: none"> Pure red cell aplasia (PRCA) Posterior reversible encephalopathy syndrome (PRES) Torsades de points Approval duration: 12 months
<p>tadalafil (Adcirca; Alyq) PAH: 20 mg tablets</p> <p><i>Tadalafil for treatment of BPH is non-formulary. If medically necessary,</i></p>	<ol style="list-style-type: none"> Ordered for an approved indication for use: <ul style="list-style-type: none"> To treat signs and symptoms of benign prostatic hyperplasia (BPH). To treat pulmonary arterial hypertension (World Health Organization group 1) to improve exercise 	<ol style="list-style-type: none"> All initial criteria continue to be met. IF patient is also prescribed macitentan (Opsumit), please redirect to Opsynvi (macitentan + tadalafil).

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
<p><i>must be requested under a non-formulary exception PA request.</i></p>	<p>ability.</p> <ol style="list-style-type: none"> 2. Patient is not concurrently on organic nitrates (i.e., isosorbide mononitrate, isosorbide dinitrate, nitroglycerin), OR sildenafil, OR Adempas (riociguat); AND 3. IF patient is also prescribed macitentan (Opsumit), please redirect to Opsynvi (macitentan + tadalafil). 4. Erectile dysfunction is not a covered indication for use. 5. Ordered for generic Adcirca (tadalafil PAH) 20 mg tablets. 6. The diagnosis of PAH is documented by right-heart catheterization with ALL of the following: <ul style="list-style-type: none"> ○ Mean pulmonary artery pressure (mPAP) > 20 mmHg, and ○ Pulmonary arterial wedge pressure (PAWP) ≤ 15 mmHg, and ○ Pulmonary vascular resistance (PVR) ≥ 3 wood units. 7. Medication ordered by a Pulmonologist, Cardiologist, or Rheumatologist. 8. Quantity Limits: 2 tablets per day. 9. Approval Duration: 12 months. 	<ol style="list-style-type: none"> 3. Patient has documented positive clinical response to tadalafil treatment. 4. Approval Duration: 12 months.
<p>tenofovir alafenamide (Vemlidy) tablets 25mg</p>	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Treatment of chronic hepatitis B virus infection in adults and pediatric patients, ≥ 6 years of age and weighing at least 25 kg, with compensated liver disease 2. Baseline test results prior to treatment start. <ul style="list-style-type: none"> • Confirmed negative HIV test result prior to starting medication. • Hepatitis Be antigen (HBeAg) status. 	<ol style="list-style-type: none"> 1. Documentation of a positive clinical response to Vemlidy therapy. 2. Patient is not a suitable candidate for entecavir. 3. Approval duration: 12 months.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> • Liver function tests. Not recommended for Child-Pugh class B or C hepatic impairment. <ol style="list-style-type: none"> 3. Patient has a history of adverse event, intolerance to or contraindication to treatment with entecavir OR meets one of the following criteria: <ul style="list-style-type: none"> • Patient age < 20 years. • Documentation of osteopenia or osteoporosis as defined by a T-score ≤ 1 and supported by clinical documentation of DEXA scan results. • Submission of medical records documenting a prior low-trauma or non-traumatic fracture. 4. In patients with renal impairment, patients who are not receiving chronic hemodialysis must have an estimated creatinine clearance > 15 ml/minute 5. Medication ordered or in consultation with an Infectious Disease specialist, Gastroenterologist, or Hepatologist. 6. Initial authorization period: 12 months. 	
teriparatide (Forteo) 620mcg/2.48ml Pen-injector	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Treatment of postmenopausal women with osteoporosis at high risk for fracture. • To increase bone mass in men with primary or hypogonadal osteoporosis at high risk of fracture. • Treatment of men and women with osteoporosis associated with sustained systemic glucocorticoid therapy (daily dose equivalent to ≥ 5 mg of prednisone) at high risk for fracture. 2. Age ≥ 18 years or documentation of closed epiphyses on X-ray. 	Osteoporosis <ol style="list-style-type: none"> 1. Patient previously met initial approval criteria. 2. Documentation supports positive response to therapy. 3. If request is for continuation of cumulative PTH analog therapy beyond 2 years, provider attestation that member remains at or has returned to having a high risk for fracture (e.g., history of osteoporotic fracture or multiple risk factors for fracture) and that the risk versus benefit of continued therapy has been reviewed with the member.

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ol style="list-style-type: none"> 3. Patient is at very high fracture risk as evidenced by one of the following: <ul style="list-style-type: none"> • Recent osteoporotic fracture within the past 12 months. • Bone mineral density (BMD) T-score at hip or spine ≤ -3.0 • BMD T-score at hip or spine ≤ -2.5 AND major osteoporotic fracture (i.e., hip, spine, forearm, wrist, humerus). 4. Patient has completed a 3-year trial of bisphosphonate therapy at up to maximally indicated doses, UNLESS one of the following: <ul style="list-style-type: none"> • All bisphosphonates are contraindicated. • Clinically adverse effects are experienced to both IV and PO formulations. • Patient has experienced a loss of- or a lack of increase in- BMD while receiving bisphosphonate therapy. • Patient experienced an osteoporotic fracture or fragility fracture while receiving bisphosphonate therapy. 5. If request is for continuation of cumulative PTH analog therapy beyond 2 years, provider attestation that member remains at or has returned to having a high risk for fracture (e.g., history of osteoporotic fracture or multiple risk factors for fracture) and that the risk versus benefit of continued therapy has been reviewed with the member. 6. Dose does not exceed 20 mcg per day (1 pen every 28 days) 7. Approval Duration: 6 months. 	<ol style="list-style-type: none"> 4. If request is for a dose increase, the new dose does not exceed 20 mcg per day (1 per per 28 days). 5. Approval duration: 12 months <p><u>Glucocorticoid-induced osteoporosis:</u></p> <ol style="list-style-type: none"> 1. Documentation supports positive response to therapy. 2. If request is for continuation of cumulative PTH analog therapy beyond 2 years, provider attestation that member remains at or has returned to having a high risk for fracture (e.g., history of osteoporotic fracture or multiple risk factors for fracture) and that the risk versus benefit of continued therapy has been reviewed with the member. 3. Approval duration: not to exceed 6 months.
tesamorelin (Egrifta SV) injection 2mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • Reduction of excess abdominal fat in HIV-infected adult patients with lipodystrophy. 2. Diagnosis of HIV-associated lipodystrophy. 	<ol style="list-style-type: none"> 1. Documentation of positive clinical response (e.g., improvement in visceral adipose tissue [VAT], decrease in waist circumference, belly appearance).

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ol style="list-style-type: none"> 3. Patient age \geq 18 years and \leq 65 years. 4. Patient meets ONE of the following: <ul style="list-style-type: none"> • If male, waist circumference is \geq 95 cm (37.4 inches) and waist-to-hip ratio is \geq 0.94; OR • If female, waist circumference is \geq 94 cm (37 inches) and waist-to-hip ratio is \geq 0.88; AND 5. Patient has been stable on antiretroviral regimen for at least 8 weeks; AND 6. Medication is prescribed by or in consultation with an endocrinologist or physician specializing in the treatment of HIV-infection. 7. Approval Duration: 6 months. 	<ol style="list-style-type: none"> 2. Approval Duration: 12 months.
tirzepatide (Mounjaro) injection 2.5mg/0.5ml, 5mg/0.5ml, 7.5mg/0.5ml, 10mg/0.5ml, 12.5mg/0.5ml, 15mg/0.5ml	<ol style="list-style-type: none"> 1. Ordered for the covered indication: <ul style="list-style-type: none"> • Treatment of adult patients with Type 2 Diabetes mellitus (T2DM). 2. Patient age \geq 18 years. 3. Patient has diagnosis of T2DM. NOTE: this product is not indicated for use in T1DM. 4. A1c or Time in Range% (TIR%) CGM-report within past 3 months. 5. A urine albumin-to-creatinine ratio (uACR) within the previous 12 months. <p><u>Treatment of Type 2 Diabetes without regard to CVD risk factors:</u> The patient has an A1c (hemoglobin A1c) of \geq 7.5 (TIR \leq 60%).</p> <p style="text-align: center;">OR</p> <p><u>Treatment of Type 2 Diabetes with CVD as defined below:</u></p> <ul style="list-style-type: none"> • Pre-treatment A1c is \geq 6.5 (TIR \leq 70%) AND 	<p>Cannot be approved for indication of weight management.</p> <ol style="list-style-type: none"> 1. Chart notes with A1c or CGM report with TIR% within previous 3 months. 2. A urine albumin-to-creatinine ratio (uACR) within the previous 12 months. 3. Documented positive clinical response defined as one of the following: <u>Baseline (pre-GLP-1) A1c was \geq 8.0 and:</u> <ul style="list-style-type: none"> • A1c has decreased by \geq 1% since onset of therapy or TIR% was \leq 55% and has increased \geq 10% or 5. A1c is \leq 7.0 at initiation dose. <u>Baseline (pre-GLP-1) A1c was \geq 6.5 but $<$ 8.0 and:</u> 6. A1c or TIR% has improved. Not eligible for renewal if A1c has increased or TIR% has decreased. <ol style="list-style-type: none"> 4. Patient has not had medical

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> • BMI \geq 27 kg/m² (documentation within previous 90 days of current height and weight); AND <p>Documentation submitted to show that the patient has at least one of the following:</p> <ul style="list-style-type: none"> • History of myocardial infarction; or • Prior stroke (ischemic or hemorrhagic); or • Symptomatic peripheral arterial disease (PAD) as evidenced by: <ul style="list-style-type: none"> ○ Intermittent claudication with ankle-brachial index (ABI) < 0.85 (at rest); or ○ Peripheral arterial revascularization procedure; or ○ Amputation due to atherosclerotic disease. <p>6. May not be concurrently using or taking ANY of the following:</p> <ul style="list-style-type: none"> • ANY other GLP1 or GLP1/GIP combination drug (e.g., Ozempic, Rybelsus, Saxenda, Soliqua, Trulicity, Victoza (liraglutide), Xultrophy, or Zepbound) • ANY DPP4i (e.g., alogliptin, Januvia (sitagliptin), Onglyza (saxagliptin) or Tradjenta (linagliptin)). • Agents for <i>severe</i> constipation: metoclopramide, Amitiza (lubiprostone), Linzess (linaclotide), Motegrity (prucalopride) or Trulance (plecanatide). <p>7. Prescriber attests that medication is prescribed in accordance with prescribing information, including screening for any black box warnings and all contraindications.</p>	<p>intervention for:</p> <p>7. Pancreatitis; or</p> <p>8. Severe gastrointestinal events. (e.g., hospitalization or new start GI motility agent).</p> <p>5. May not be concurrently using ANY of the following:</p> <ul style="list-style-type: none"> • ANY other GLP1 or GLP1/GIP combination drug (e.g., Ozempic, Rybelsus, Saxenda, Soliqua, Trulicity, Victoza, Xultrophy or Zepbound) • ANY DPP4i (e.g., alogliptin, Januvia (sitagliptin), Onglyza (saxagliptin), or Tradjenta (linagliptin)). • Agents for <i>severe</i> constipation: metoclopramide, Amitiza (lubiprostone), Linzess (linaclotide), Motegrity (prucalopride) or Trulance (plecanatide). <p>5. PBM claims data shows consistent adherence as shown by no instance of a drug-free interval greater than 2 months at which time the patient would need to satisfy the initial criteria.</p> <p>6. Approval Duration: up to 12 months.</p>

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>8. May not be approved for patients with:</p> <ul style="list-style-type: none"> • Any personal or family history of medullary thyroid carcinoma (MTC) or multiple endocrine neoplasia syndrome type 2 (MEN 2). • Current pregnancy • History of confirmed pancreatitis <p>9. Starter doses are limited and require dose escalation. Mounjaro 2.5 mg is a starter dose and is limited to one, 28-day supply and then must be dose escalated to 5 mg per week dose UNLESS A1c ≤ 7.0 or TIR ≥ 65% on 2.5 mg dose.</p> <p>10. Cannot be approved for indication of weight management.</p> <p>11. Maximum Approval Duration: up to 12 months</p>	
<p>tivozanib (Fotivda) capsules 0.89mg, 1.34mg</p>	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • the treatment of adult patients with relapsed or refractory advanced renal cell carcinoma (RCC) following two or more prior systemic therapies. 2. Patient has relapsed or Stage IV disease; AND 3. Patient has tried at least two other systemic regimens (i.e. Inlyta + Keytruda; Cabometyx + Opdivo; Lenvima + Keytruda; Yervoy + Opdivo, sunitinib, pazopaniv, or Lenvima + everolimus. 4. Medication order by Hematology/oncology. 5. Approval Duration: 12 months. 	<ol style="list-style-type: none"> 1. Patient does not show evidence of disease progression while on Fotivda therapy. 2. Approval Duration: 12 months.
<p>tramadol hydrochloride extended release (Ultram) capsules (biphasic release) 100mg, 150mg, 200mg, 300mg</p> <p>Tablets 100mg, 200mg, 300mg</p>	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • in adults for the management of pain severe enough to require an opioid analgesic and for which alternative treatments are inadequate. 2. Completion of the opioid prior authorization form. 3. Submission of supporting clinical documentation for last office visit, dated within the previous 3 months. 	<p>All long-acting opioids require Prior Authorization (PA). The PA form can be accessed using the following link: OPIOID PRIOR AUTHORIZATION FORM-DC</p> <p>Limitations of Use:</p>

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
Tablets (biphasic release) 100mg, 200mg, 300m	<ol style="list-style-type: none"> 4. Opioid naïve adult patients (no opioid use within the previous 30-days) are limited to a 7-day supply with an MME ≤ 50 MME per day, patients under age 18 are limited to a 3-day supply. 5. Maximum approval duration is 6 months but may be reduced or denied based on any of the criteria as outlined in Pharmacy Policy 219.DC: Opioid Prescription Prior Authorization. 	Not indicated as an as-needed (prn) analgesic.
treprostinil (Orenitram) 0.125 mg, 0.25 mg, 1 mg, 2.5 mg, 5 mg tablets	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • To delay disease progression and improve exercise capacity in patients with WHO functional class II-III symptoms and etiologies of idiopathic or heritable PAH or PAH associated with connective tissue disease. 2. Patient aged ≥ 18 years. 3. Patient has a diagnosis of World Health Organization (WHO) Group I pulmonary arterial hypertension (PAH); AND 4. Documentation is submitted showing that the patient has had a right heart catheterization that confirms the diagnosis of WHO Group I PAH; AND 5. Patient meets one of the following conditions: <ul style="list-style-type: none"> • Patient has tried TWO oral therapies for PAH from two of the three different categories (either alone or in combination) of each for ≥ 60 days: <ul style="list-style-type: none"> ○ One phosphodiesterase type 5 (PDE5) inhibitor (e.g. sildenafil or tadalafil); or ○ One endothelin receptor antagonist (ERA) (e.g. bosentan, ambrisentan or Opsumit [macitentan]), or Adempas (riociguat); OR 	<ol style="list-style-type: none"> 1. Patient has a diagnosis of World Health Organization (WHO) Group I pulmonary arterial hypertension (PAH); AND 2. Patient has had a right heart catheterization; AND 3. The results of the right heart catheterization confirm the diagnosis of WHO Group I PAH; AND 4. The patient is experiencing a positive clinical response to treatment with Orenitram as evidenced by any of the following: reduced pulmonary vascular resistance and/or pressure, improved symptoms, and/or improved patient activity. 5. Not prescribed concurrently with Upravi (selexipag), inhaled prostacyclin products (e.g. Tyvaso [Treprostinil], Tyvaso DPI, Ventavis [iloprost], epoprostenol). 6. Orenitram is prescribed by, or in consultation with a cardiologist or pulmonologist. 7. Approval Duration: 12 months.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> ○ Patient has history of treatment with one PAH prostacyclin therapy or a prostacyclin receptor agonist 6. Patient does not have severe hepatic impairment (Child Pugh Class C). 7. Not prescribed concurrently with Uptravi (selexipag), inhaled prostacyclin products (e.g. Tyvaso [Treprostini], Tyvaso DPI, Ventavis [iloprost], epoprostenol). 8. Prescribed by or in consultation with a cardiologist or pulmonologist. 9. Quantity Limit: 2 tablets per day. Use appropriate tablet strength to reach desired total daily dose. 10. Approval Duration: 12 months. 	
triptorelin (Trelstar) intramuscular injection 3.75 mg; 11.25 mg; 22. 5 mg	1. Ordered for an approved indication for use: <ul style="list-style-type: none"> ● Palliative treatment of advanced prostate cancer ● Preservation of ovarian function ● Breast cancer (ovarian suppression) ● Gender affirming care. 2. Prostate Cancer: <ul style="list-style-type: none"> ● Prescribed by an oncologist. Preservation of ovarian function: <ul style="list-style-type: none"> ● Patient is premenopausal and undergoing chemotherapy. Breast cancer: <ul style="list-style-type: none"> ● Patient is premenopausal with hormone-receptor positive breast cancer at high-risk for recurrence using in combination with endocrine therapy. Gender affirming care: <ul style="list-style-type: none"> ● Patient has diagnosis of gender dysphoria and meets MDH regulatory requirements for care. ● Patient has reached Tanner stage ≥ 2 of puberty. 	1. Prostate Cancer: <ul style="list-style-type: none"> ● Patient is experiencing clinical benefit (e.g., serum testosterone < 50 ng/dl) ● Patient has not experienced unacceptable toxicity. Preservation of ovarian function: <ul style="list-style-type: none"> ● Patient meets all initial criteria. Breast cancer: <ul style="list-style-type: none"> ● Patient was premenopausal at diagnosis and is still undergoing treatment with endocrine therapy. ● Total treatment with triptorelin does not exceed 5 years. Gender affirming care: <ul style="list-style-type: none"> ● Patient has reached Tanner stage ≥ 2 of puberty. 2. Approval Durations:

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>3. Approval Durations:</p> <ul style="list-style-type: none"> • Prostate Cancer: 12 months • Preservation of ovarian function: 3 months • Breast cancer (ovarian suppression): 12 months • Gender affirming care: 12 months 	<ul style="list-style-type: none"> • Prostate Cancer: 12 months • Preservation of ovarian function: up to 12 months <i>**providing that cumulative treatment course is < 5 years.</i> • Breast cancer: 12 months • Gender affirming care: 12 months
ubrogepant (Ubrelyv) tablets 50mg, 100mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • the acute treatment of migraine with or without aura in adults. 2. Patient age ≥ 18 years. 3. Member must have tried and failed NSAIDs and at least two formulary triptans or have a contraindication to taking both classes of medications. *examples of contraindications include: a history of coronary artery disease, cardiac accessory pathway disorders, history of stroke or TIA, or hemiplegic or basilar migraine, peripheral vascular disease, ischemic bowel disease, uncontrolled hypertension, or severe hepatic impairment. 4. Quantity limited to 16 doses per 30 days, 200 mg max daily dose. 5. Approval Duration: 12 months. 	<ol style="list-style-type: none"> 1. Meets all initial clinical criteria. 2. Documentation of positive clinical response to treatment. 3. Quantity limited to 16 doses per 30 days, 200 mg max daily dose. 4. Approval Duration: 12 months.
ustekinumab (Stelara) Injection, 45 mg; 90 mg	<ol style="list-style-type: none"> 1. The criteria for Stelara are indication specific. Please review criteria for the patient-specific diagnosis. 2. Stelara induction therapy requires Prior Authorization and must meet the prior authorization criteria below. 3. Patient has been screened for Hepatitis B and Tuberculosis prior to initiation of therapy. 	<p>ALL INDICATIONS:</p> <ol style="list-style-type: none"> 1. Documented positive clinical response. 2. Patient is not receiving in combination with any other targeted immunomodulator (e.g., etanercept, certolizumab, golimumab, abatacept, adalimumab, Risankizumab, guselkumab, secukinumab, ixekizumab, brodalumab,

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>4. Patient is not receiving in combination with any other targeted immunomodulator (e.g., etanercept, certolizumab, golimumab, abatacept, adalimumab, Risankizumab, gueslkumab, secukinumab, ixekizumab, brodalumab, tildrakizumab, rofacitinib, baricitinib, upadacitinib, apremilast, or similar).</p> <p>Hidradenitis suppurative: <u>excluded</u> from coverage; off-label indication. Note: Humira (or biosimilars) is first line therapy. Remicade (infliximab) is the MFC recommended alternate.</p> <p>Crohn's disease:</p> <ul style="list-style-type: none"> • Diagnosis of moderately to severely active Crohn's disease • Patient is currently on Stelara therapy for moderately to severely active Crohn's disease as documented by claims history or submission of medical records. • Must have trialed and failed therapy with adalimumab, this includes patients who have failed infliximab. (1A recommendation from AGA Practice Guidelines 2021). • Approved dose: 90 mg/ml • Approval Duration: 12 months <p>Plaque psoriasis:</p> <ul style="list-style-type: none"> • Diagnosis of moderate to severe plaque psoriasis • ≥ 3% body surface area involvement, palmoplantar, facial, genital involvement, or severe scalp psoriasis; AND • History of failure to one of the following topical therapies: <ul style="list-style-type: none"> ○ Corticosteroids ○ Vitamin D analogs (calcitriol, calcipotriene) 	<p>tildrakizumab, rofacitinib, baricitinib, upadacitinib, apremilast, or similar).</p> <p>3. Approval Duration: 12 months.</p>

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ul style="list-style-type: none"> ○ Tacrolimus or pimecrolimus. ● History of failure to a 3-month trial of methotrexate at maximally indicated dose, unless contraindicated or clinically significant adverse effects are experienced. The trial must be documented in chart notes with date and duration of trial, OR ● Patient has been previously treated with a targeted immunomodulator indicated for the treatment of plaque psoriasis as documented by claims history or submission of medical records that include drug name, date, and duration of therapy. (e.g., adalimumab, certolizumab, apremilast, Risankizumab, gueslkumab or similar). ● Must be prescribed by or in consultation with a dermatologist. ● Patients requesting initial authorization who were established on therapy via the receipt of a manufacturer supplied sample at no cost in the prescriber’s office or any form of assistance from the Janssen sponsored CarePath Savings program shall be required to meet initial authorization criteria as if the patient were new to therapy. ● Approved dose: 45 mg/ml for weight ≤ 100 kg ● Approved dose: 90 mg/ml for weight > 100 kg ● Approval duration: 12 months. <p>Psoriatic arthritis:</p> <ul style="list-style-type: none"> ● Diagnosis of active psoriatic arthritis; AND ● History of failure to a 3-month trial of methotrexate at maximally indicated dose, unless contraindicated or clinically significant adverse 	

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<p>effects are experienced. The trial must be documented in chart notes with date and duration of trial, OR</p> <ul style="list-style-type: none"> • Patient has been previously treated with a targeted immunomodulator indicated for the treatment of plaque psoriasis as documented by claims history or submission of medical records that include drug name, date, and duration of therapy. (e.g., adalimumab, certolizumab, apremilast, golimumab, guselkumab, tofacitinib, upadacitinib, or similar). • Patients requesting initial authorization who were established on therapy via the receipt of a manufacturer supplied sample at no cost in the prescriber’s office or any form of assistance from the Janssen sponsored CarePath Savings program shall be required to meet initial authorization criteria as if the patient were new to therapy. • Prescribed by or in consultation with a rheumatologist or dermatologist. • Approved dose: 45 mg/ml for weight ≤ 100 kg • Approved dose: 90 mg/ml for weight > 100 kg • Approval duration: 12 months <p><u>Ulcerative colitis, moderate to severe:</u></p> <ul style="list-style-type: none"> • Must show treatment failure or contraindication to first-line therapies: Remicade (infliximab) or Entyvio (vedolizumab). • Prescribed by or in consultation with a gastroenterologist. • Approved dose: 90 mg/ml • Approval Duration: 12 months 	
valbenazine (Ingrezza) capsules	1. Ordered for an approved indication for use:	1. Prescriber attestation of continued

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
40 mg, 60 mg, 80 mg	<ul style="list-style-type: none"> • Chorea associated with Huntington’s disease. (HD) • Tardive dyskinesia (TD) in adults. <ol style="list-style-type: none"> 2. Patient age ≥ 18 years. 3. Patient is not receiving other VMAT2 inhibitors (tetrabenazine or deutetrabenazine), MAOIs or reserpine. 4. Patient does not have hepatic impairment. 5. Tardive dyskinesia: <ul style="list-style-type: none"> • AIMS score sheet along with the progress note must be provided for initial and renewal PA requests. 6. Huntington’s disease: <ul style="list-style-type: none"> • Description of functional impairment, including Total Maximal Chorea (TMC) score sheet along with progress notes must be provided for both initial and renewal PA requests. 7. Patient must not be suicidal or have untreated/inadequately treated depression. 8. Patient is not prescribed to concurrently use multiple strengths of Ingrezza. 9. Quantity Limit: 1 capsule per day. 10. Approval Duration: 12 months 	<p>clinical benefit and subsequent evaluation and monitoring performed.</p> <ol style="list-style-type: none"> 2. TD: AIMS score must show improvement over initial score. 3. HD: TMC score must show improvement over the initial score and functional impairment must show improvement from baseline. 4. All initial criteria must be met. 5. Approval Duration: 12 months
vigabatrin (Sabril; Vigadrone) 500 mg powder pack	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • treatment of Refractory Complex Partial Seizures as adjunctive therapy in patients ≥ 2 years of age who have responded inadequately to several alternative treatments. • Infantile Spasms - monotherapy in infants 1 month to 2 years of age for whom the potential benefits outweigh the potential risk of vision loss. 	<p>Limitations of Use: Vigabatrin is not indicated as a first line agent.</p>

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	2. Medication prescribed by a provider registered in the vigabatrin REMS program. 3. Medication prescribed by a Neurologist.	
viloxazine extended release (Qelbree) capsules 100mg, 150mg, 200mg STEP THERAPY	1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • treatment of Attention Deficit Hyperactivity Disorder (ADHD) in patients ≥ 6 years of age. 2. History of a ≥4-week trial of, contraindication to, or intolerance of both of the following stimulant medications: <ul style="list-style-type: none"> • a methylphenidate class stimulant (e.g. generic Concerta) • an amphetamine class stimulant (e.g. generic Adderall XR) • <u>OR</u> history of a substance use disorder or concern for potential misuse or diversion of stimulant medication <u>AND</u> 3. History of a ≥4-week trial of, contraindication to, or intolerance of both of the following non-stimulant medications: <ul style="list-style-type: none"> • guanfacine ER (generic Intuniv) • atomoxetine (generic Straterra) 4. Daily dose does not exceed 400 mg per day for patients 11 years of age or younger; 600 mg per day max for ages ≥12 years.	
viltolarsen (Viltepsa) 250 mg/5ml solution	1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • treatment of Duchenne muscular dystrophy (DMD) in patients who have a confirmed mutation of the DMD gene that is amenable to exon 53 skipping. 2. Genetic testing must confirm patient’s DMD gene is amenable to exon 53 skipping.	1. Documentation and provider attestation of continued benefit, including respiratory status assessment, without adverse effects. 2. Not receiving another antisense therapy or gene therapy.

Generic Medication (Brand Name) Bolded name indicates whether Brand or Generic is Formulary	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	<ol style="list-style-type: none"> 3. Current patient weight, including date weight was obtained and within 30 days of requested date. 4. Baseline renal function test (GFR) and Urine protein-to-creatinine ratio prior to starting treatment. 5. Documented baseline function testing using a tool to demonstrate physical functions, including, but not limited to: Brooke Upper Extremity Scale, Baseline 6-minute walk test, Pediatric Evaluation of Disability Inventory. 6. Stable dose of glucocorticoid for at least 3 months. 7. Confirmation that drug continues to carry FDA-approval for indication. 8. Prescribed by a neurologist with expertise in treatment of DMD. 9. Approval duration: 3 months 	<ol style="list-style-type: none"> 3. Verification that drug continues to carry FDA-approval for indication. <ul style="list-style-type: none"> • Approval duration: 3 months
voclosporin (Lupkynis) capsule 7.9mg	<ol style="list-style-type: none"> 1. Ordered for an approved indication for use: <ul style="list-style-type: none"> • in combination with a background immunosuppressive therapy regimen for the treatment of adult patients with active lupus nephritis. (mycophenolate mofetil and corticosteroids). 2. Patient age \geq 18 years. 3. Not taking concurrently with cyclophosphamide. 4. Prescriber specialty: immunologist, nephrologist, rheumatologist, or provider experienced in treatment of lupus nephritis. 5. Prescriber attestation that all baseline evaluations have been done, and not contraindications to use are present (strong 3A4 inhibitor contraindicated, live vaccines, pregnancy/breastfeeding negative, assessment of renal function). 6. Quantity Limit: 6 tablets per day (23.7 mg twice daily). 	<ol style="list-style-type: none"> 1. All initial criteria continue to be met. 2. Documentation provided or attestation of therapeutic benefit. 3. Approval Duration: 6 months.

Generic Medication (Brand Name) <small>Bolded name indicates whether Brand or Generic is Formulary</small>	Approval Criteria & Submission Requirements	Additional Considerations & Renewal Criteria
	7. Approval Duration: 6 months	